

PRECISE ASYMPTOTICS OF THE RICCI FLOW NECKPINCH

SIGURD ANGENENT & DAN KNOFF

CONTENTS

1. Introduction	2
1.1. Antecedents	2
1.2. Overview	3
1.3. A diameter bound	4
2. Rigorous asymptotics of the $\mathrm{SO}(\mathbb{R}^{n+1})$ invariant neckpinch	5
2.1. The blown-up radius and the linearized equation	5
2.2. Pointwise estimates for u and v	7
2.3. Transition to commuting variables	8
2.4. The linearized equation	10
2.5. Eigenfunction decomposition of v	10
2.6. Easy estimates for a_0 and a_2	11
2.7. Decay of a_0	11
2.8. Decay of W	12
2.9. Derivative estimates for W	13
2.10. A bound for $\ \sigma W\ $	14
2.11. Pointwise estimates for W	14
2.12. Pointwise estimate for W_σ	15
2.13. An equation for a_2	15
2.14. Dichotomy for the decay of $a_2(\tau)$	18
2.15. Pinching time	19
2.16. Initial data	20
2.17. The dominant asymptotic profile	22
2.18. The intermediate region	24
2.19. Convergence in the outer region	28
2.20. Asymptotics in the outer region	31
3. Formal asymptotics for general neckpinches	31
3.1. The cylinder soliton	32
3.2. The dilated Ricci flow	33
3.3. First and second order variation formulas	34
3.4. Arbitrary perturbations of a stationary solution	35
3.5. The linearization at the cylinder soliton	36
3.6. An upper bound for the spectrum of the linearization	37
3.7. A decomposition into invariant subspaces	38
3.8. The spectrum of the linearization	39
3.9. Our <i>Ansatz</i>	41

First author partially supported by NSF grant DMS-0101124. Second author partially supported by NSF grants DMS-0511184, DMS-0505920, and a University of Texas Summer Research Assignment.

3.10.	The inner layer $ x = o(\sqrt{\tau})$	43
3.11.	The intermediate layer $ x = \mathcal{O}(\sqrt{\tau})$	46
3.12.	The outer layer $ x = o(1/\sqrt{T-t})$	46
	References	47

1. INTRODUCTION

1.1. Antecedents. In virtually all known applications of the Ricci flow, it is valuable to have a good understanding of singularity formation. Heuristically, there are at least three reasons for this. The first is that one expects finite-time singularities to form for a broad spectrum of initial data. Indeed, such singularities are inevitable if the scalar curvature is strictly positive. The second reason is that one expects the geometry of a solution to resemble a standard model (for example, a self-similar solution) in a space-time neighborhood of a developing singularity. The third reason is that having a sufficiently detailed picture of a developing singularity facilitates the geometric-topological surgeries by which Ricci flow decomposes a given manifold.

Whenever a compact solution $(M^n, g(\cdot))$ of Ricci flow encounters a singularity at time $T < \infty$, standard short-time existence results imply that

$$\lim_{t \nearrow T} \max_{x \in M^n} |\text{Rm}(x, t)| = \infty.$$

Recent results of N. Sesum [28] allow one to replace this condition by

$$\limsup_{t \nearrow T} \max_{x \in M^n} |\text{Rc}(x, t)| = \infty.$$

The most interesting cases are those where a *local singularity* forms, that is, where there exists an open set $\Omega \subset M^n$ such that

$$\sup_{\Omega \times [0, T)} |\text{Rc}(x, t)| < \infty.$$

The first rigorous constructions of local singularities were done by M. Simon [29]. Here the manifold is a noncompact warped product $\mathbb{R} \times_f S^n$, and a supersolution of the Ricci flow PDE is used to prove that f vanishes in finite time on a proper subset of $\mathbb{R}$. A second class of examples was constructed in [12]. Here the manifold is a complex line bundle $\mathbb{C} \hookrightarrow \mathbb{L}_{-k}^n \rightarrow \mathbb{CP}^{n-1}$ with $1 \leq k \leq n-1$. As the singularity forms, the flow performs an algebraic-geometric blow-down of the $\mathbb{CP}^{n-1}$, while the rest of the manifold converges locally smoothly to a metric cone on $(\mathbb{C}^n \setminus \{0\})/\mathbb{Z}_k$.

As G. Perelman writes, “The most natural way of forming a singularity in finite time is by pinching an (almost) round cylindrical neck.” [27] This is the kind of local singularity that we analyze in this paper. It is a continuation of our earlier work [3], where we gave the first rigorous examples of local singularity formation on compact manifolds by constructing neckpinches for rotationally-symmetric metrics on S^{n+1} . In that paper, we obtained local *a priori* estimates for the space-time scales at which a developing neckpinch singularity resembles the self-similarly-shrinking cylinder soliton. The present paper shows that those estimates are sharp, and gives precise asymptotics for neckpinch formation.

When considering a local singularity, two natural and important questions arise. (1) What is the nature of the set of points in space on which the metric becomes singular? (2) What is the asymptotic behavior of the solution near this set as the singularity time is approached? In the past two decades, a rich literature of both rigorous and formal matched asymptotics has developed for analyzing the local behavior of singular solutions of nonlinear PDE such as $u_t = \Delta(\log u)$ or $u_t = \Delta u + F(u)$, where $F(u) = u^p$ or $F(u) = e^u$. Some of the many noteworthy results of this type are [32], [26], [14], [18], [15, 16, 17], [20, 21, 22], [25], and [13]. A few results specific to geometric evolution equations such as mean curvature flow and harmonic map flow are [1], [4, 5], [31], [2], and [8].

1.2. Overview. This paper is divided into two main parts.

Rigorous asymptotics. In Section 2, we derive rigorous asymptotics for rotationally symmetric neckpinches. The set-up is as follows. Let g_{can} denote the round metric of radius 1 on S^n . Then any $\text{SO}(n+1)$ -invariant metric on S^{n+1} can be written as

$$g = \varphi^2 dx^2 + \psi^2 g_{\text{can}}$$

on $(-1, 1) \times S^n$, which may be naturally identified with the sphere S^{n+1} with its north and south poles removed. The quantity $\psi(x, t) > 0$ may thus be regarded as the radius of the hypersurface $\{x\} \times S^n$ at time t . It is natural to write geometric quantities related to g in terms of the distance $s(x) = \int_0^x \varphi(\hat{x}) d\hat{x}$ from the equator. Then one can write the metric in the nicer form of a warped product

$$g = ds^2 + \psi^2 g_{\text{can}},$$

bearing in mind that s is ultimately a function of both x and t . (Note that one encounters the commutator $[\partial_t, \partial_s] = -n\psi^{-1}\psi_{ss}\partial_s$ when taking derivatives with x held constant.)

In [3], we established neckpinching for a class of data essentially described by three conditions: (1) the initial metric should have positive scalar curvature on all of S^{n+1} and positive Ricci curvature on the polar caps (in the terminology of [3]); (2) its sectional curvature should be positive on planes tangential to the spheres $\{x\} \times S^n$; and (3) it should be ‘sufficiently pinched’, i.e. the minimum radius should be sufficiently small relative to the maximum radius. In this paper, we impose an additional hypothesis: (4) the initial metric should be reflection symmetric, i.e. $\psi(s, 0) \equiv \psi(-s, 0)$. Under these hypotheses, the results we obtain in Section 2 (combined with results from the predecessor [3] to this paper) imply the the solution obeys a precise asymptotic profile.

To describe this profile requires some additional notation. If $T < \infty$ is the singularity time, let $u = \psi/\sqrt{2(n-1)(T-t)}$ be the blown-up radius defined below in (2); let $\sigma = s/\sqrt{T-t}$ be the rescaled distance to the neck defined in (3); and let $\tau = -\log(T-t)$ denote the rescaled time variable defined in (5). Note that (σ, τ) are self-similar coordinates with respect to the blown-up flow. We summarize our results as

Theorem 1. *For an open set of initial metrics symmetric with respect to reflection and rotation, the solution $(S^{n+1}, g(t))$ of Ricci flow becomes singular at $T < \infty$. Its diameter remains bounded for all $t \in [0, T)$. The singularity occurs only on the hypersurface $\{0\} \times S^n$. The solution satisfies the following asymptotic profile.*

Inner region: on any interval $|\sigma| \leq A$, one has

$$u(\sigma, \tau) = 1 + \frac{\sigma^2 - 2}{8\tau} + o\left(\frac{1}{\tau}\right) \quad \text{uniformly as } \tau \rightarrow \infty.$$

Intermediate region: on any interval $A \leq |\sigma| \leq B\sqrt{\tau}$, one has

$$u(\sigma, \tau) = \sqrt{1 + (1 + o(1)) \frac{\sigma^2}{4\tau}} \quad \text{uniformly as } \tau \rightarrow \infty.$$

Outer region: for any $\varepsilon > 0$, there exist $C < \infty$ and $\bar{t} < T$ such that

$$\left(\frac{1}{2}\sqrt{n-1} - \varepsilon\right) \frac{s}{\sqrt{\log(1/s)}} \leq \psi(s, t) \leq \left(\frac{1}{2}\sqrt{n-1} + \varepsilon\right) \frac{s}{\sqrt{\log(1/s)}}$$

for all $|\sigma| \geq C\sqrt{\tau}$ and $t \in (\bar{t}, T)$.

Formal asymptotics. In Section 3, we derive formal matched asymptotics for fully general neckpinches, without any symmetry assumptions whatsoever. These are analogous to the formal asymptotics for the $m = 2$ (presumably typical) case of MCF singularities considered in [5]. Our method here is to study arbitrary perturbations of the self-similar cylinder soliton. That is, we consider $\tilde{g} = g + h$, where

$$g = dx^2 + 2(n-1)(T-t)g_{\text{can}}$$

is the cylinder soliton on $\mathbb{R} \times S^n$, and h is an arbitrary (small) $(2, 0)$ -tensor. To accomplish our analysis, we modify the Ricci flow so that the soliton becomes a fixed point of a related strictly parabolic flow. Because the linearization of that flow has a null eigenvalue, one expects to find a center manifold. We proceed by carrying out a quadratic variational analysis in order formally to compute the dynamics of the flow on that center manifold.

The computations involved are extensive, but the conclusion we obtain is quite satisfying: the formal matched asymptotics suggest that the behavior analyzed rigorously in Section 2 should indeed be stable for fully general Ricci flow neckpinches.

1.3. A diameter bound. We conclude this introduction with an observation that follows directly from the estimates obtained in [3]. This result confirms an expectation of G. Perelman [19].

Lemma 2. *Let $(S^{n+1}, g(t))$ be any $\text{SO}(n+1)$ -invariant solution of Ricci flow such that $g(0)$ has positive scalar curvature and positive sectional curvature on planes tangential to the spheres $\{x\} \times S^n$.*

Assume the metric $g(t)$ has at least two bumps for all $t < T$ (in the language of [3]). Let $x = a(t)$ and $x = b(t)$ be the locations of the left- and right-most bumps, and assume that for all $t < T$ one has $\psi(a(t), t) \geq c$, $\psi(b(t), t) \geq c$ for some constant $c > 0$.

If $g(t)$ becomes singular at $T < \infty$, then $\text{diam}(S^{n+1}, g(t))$ remains bounded as $t \nearrow T$.

Proof. By Proposition 5.4 of [3], the limit profile $\psi(\cdot, T)$ exists. Let $a(t) < b(t)$ in $(-1, 1)$ denote the left-most and right-most bumps, respectively. Then $a(t) \rightarrow a(T)$ and $b(t) \rightarrow b(T)$. By Lemma 5.6, the Ricci curvature is positive (so distances are decreasing) on $(-1, a(t)]$ and $[b(t), 1)$. Hence it will suffice to bound $\text{dist}_{g(t)}(x_1, x_2)$ for arbitrary $x_1 < x_2$ in $(a(T) - \varepsilon, b(T) + \varepsilon) \subset (-1, 1)$.

Equations (5) and (11) of [3] imply that

$$\begin{aligned} \frac{d}{dt} \text{dist}_{g(t)}(x_1, x_2) &= \frac{d}{dt} \int_{x_1}^{x_2} \varphi(x, t) dx \\ &= n \int_{s(x_1)}^{s(x_2)} \frac{\psi_{ss}}{\psi} ds \\ &= n \left\{ \frac{\psi_s}{\psi} \Big|_{s(x_1)}^{s(x_2)} + \int_{s(x_1)}^{s(x_2)} \left(\frac{\psi_s}{\psi} \right)^2 ds \right\}. \end{aligned}$$

Proposition 5.1 of [3] bounds ψ_s uniformly, while Lemma 5.5 shows that the number of bumps and necks is nonincreasing in time. It follows that

$$\int_{s(x_1)}^{s(x_2)} \left(\frac{\psi_s}{\psi} \right)^2 ds \leq C \int_{s(x_1)}^{s(x_2)} \frac{|\psi_s|}{\psi^2} ds \leq C \left[\frac{1}{\psi_{\min}(t)} - \frac{1}{\psi_{\max}(t)} \right] \leq \frac{C}{\psi_{\min}(t)}.$$

Hence Lemma 6.1 lets us conclude that

$$\left| \frac{d}{dt} \text{dist}_{g(t)}(x_1, x_2) \right| \leq \frac{C}{\psi_{\min}(t)} \leq \frac{C}{\sqrt{T-t}},$$

which is integrable. $\square$

Recalling Lemmata 7.1 and 10.1 of [3], one immediately obtains

Proposition 3. *Assume that $(S^{n+1}, g(t))$ satisfies the hypotheses of Lemma 2 and is reflection-symmetric. Then the final-time profile $\psi(\cdot, T)$ has $\psi(x, T) > 0$ for all $x \neq 0$. Because $|\text{Rm}| \leq C/\psi^2$, this implies that the singularity occurs only on the hypersurface $\{0\} \times S^n$.*

Diameter estimates for arbitrary finite-time Ricci flow singularities are studied from another point of view in [30]. Lower bounds for diameter are known for certain topologies; see [24].

2. RIGOROUS ASYMPTOTICS OF THE $\text{SO}(\mathbb{R}^{n+1})$ INVARIANT NECKPINCH

2.1. The blown-up radius and the linearized equation. The radius ψ satisfies

$$(1) \quad \psi_t = \psi_{ss} - (n-1) \frac{1 - \psi_s^2}{\psi}.$$

We will assume that the solution is defined on some time interval $t_0 \leq t < T$, where the initial time t_0 need not be 0. In order to construct examples of solutions that follow our precise asymptotics, we will at some point actually have to assume that $T - t_0$ is “sufficiently small”.

We consider the blown-up radius

$$(2) \quad u \doteq \frac{\psi}{R_n(t)}, \quad \text{where} \quad R_n(t) \doteq \sqrt{2(n-1)(T-t)},$$

in which T may or may not be the blow-up time. We also introduce the rescaled distance to the neck

$$(3) \quad \sigma \doteq \frac{s}{\sqrt{T-t}}.$$

(Here the absence of a factor $n-1$ in $\sqrt{T-t}$ is intentional and will result in the numerically simplest equations later on.) Then we have

$$\psi_s = \sqrt{2(n-1)} u_\sigma, \quad \psi_{ss} = \sqrt{2(n-1)} \frac{u_{\sigma\sigma}}{\sqrt{T-t}},$$

and

$$\psi_t = \sqrt{2(n-1)(T-t)} u_t - \frac{\sqrt{2(n-1)}}{2\sqrt{T-t}} u.$$

This leads to the following evolution equation for u ,

$$(4) \quad u_\tau = u_{\sigma\sigma} + \frac{1}{2} \left(u - \frac{1}{u} \right) + (n-1) \frac{u_\sigma^2}{u},$$

where

$$(5) \quad \tau \doteq -\log(T-t).$$

The blown-up time variable τ takes values in the interval

$$\tau_0 \leq \tau < \infty, \quad \text{where } \tau_0 \doteq -\log(T-t_0).$$

Without loss of generality we will assume that $\tau_0 \geq 1$.

Near the neck, u will be close to the self-similar radius $u \approx 1$. So to linearize, we set

$$(6) \quad u = 1 + v$$

and observe that v must satisfy

$$(7) \quad v_\tau = v_{\sigma\sigma} + v + \frac{(n-1)v_\sigma^2 - \frac{1}{2}v^2}{1+v}.$$

Here the partial derivatives $\partial/\partial\sigma$ and $\partial/\partial\tau$ do not commute. One has instead

$$[\partial_t, \partial_s] = -n \frac{\psi_{ss}}{\psi} \partial_s,$$

and thus

$$(8) \quad [\partial_\tau, \partial_\sigma] = \left[(T-t)\partial_t, \sqrt{T-t}\partial_s \right] = - \left\{ \frac{1}{2} + n \frac{u_{\sigma\sigma}}{u} \right\} \partial_\sigma.$$

2.2. Pointwise estimates for u and v . In [3], we showed that on solutions of the Ricci flow whose scalar curvature is nonnegative, the quantity

$$F = \frac{K}{L} \left\{ 2 + \log \frac{L}{L_{\min}(0)} \right\}$$

satisfies a maximum principle. Here

$$K = \frac{\psi_{ss}}{\psi} \text{ and } L = \frac{1 - \psi_s^2}{\psi^2},$$

and

$$L_{\min}(0) = \inf \left. \frac{1 - \psi_s^2}{\psi^2} \right|_{t=0}.$$

The maximum principle implies that $\sup F$ does not increase when it is above $n - 1$.

Let us assume throughout this paper that the initial metric has nonnegative scalar curvature, and that

$$\sup F \leq F^*$$

for some fixed constant $F^* \geq n - 1$.

When one writes the quantity F in terms of ψ , ψ_s , and ψ_{ss} , the estimate $F \leq F^*$ implies the following upper bounds for ψ , ψ_s , and ψ_{ss} in terms of the radius of the neck $\psi(0, t)$.

Lemma 4. *There exist constants $\delta > 0$ and $C < \infty$ that only depend on F^* and the dimension n such that*

$$(9) \quad 1 - \frac{C}{\tau} \leq u \leq 1 + C \frac{1 + \sigma^2}{\tau} \text{ for } |\sigma| \leq 2\sqrt{\tau},$$

and

$$(10) \quad 1 - \frac{C}{\tau} \leq u \leq C \frac{|\sigma|}{\sqrt{\tau}} \sqrt{\log \frac{|\sigma|}{\sqrt{\tau}}} \text{ for } 2\sqrt{\tau} \leq |\sigma| \leq e^{\delta\tau}.$$

For the derivative u_σ , we have

$$(11) \quad |u_\sigma| \leq C \frac{1 + |\sigma|}{\tau} \text{ for } |\sigma| \leq 2\sqrt{\tau},$$

$$(12) \quad |u_\sigma| \leq \frac{C}{\sqrt{\tau}} \sqrt{\log \frac{|\sigma|}{\sqrt{\tau}}} \text{ for } 2\sqrt{\tau} \leq |\sigma| \leq e^{\delta\tau}.$$

In both cases, these estimates imply that $|u_\sigma| \leq C\delta$ at all points with $|\sigma| \leq e^{\delta\tau}$. After decreasing δ , if necessary, we may assume that $C\delta$ is as small as we like. In particular, in the region we are considering, we always may assume that

$$(13) \quad |u_\sigma| \leq \frac{1}{2}.$$

Since $v = u - 1$, we immediately get estimates for v and v_σ . Those for v_σ are of course the same as for u_σ . For v , we have

$$(14) \quad \begin{cases} -\frac{C}{\tau} \leq v \leq C \frac{1 + \sigma^2}{\tau} \text{ for } |\sigma| \leq 2\sqrt{\tau}, \\ -\frac{C}{\tau} \leq v \leq C \frac{|\sigma|}{\sqrt{\tau}} \sqrt{\log \frac{|\sigma|}{\sqrt{\tau}}} \text{ for } 2\sqrt{\tau} \leq |\sigma| \leq e^{\delta\tau}, \end{cases}$$

in which C is again a constant depending only on F^* and n .

Proof of Lemma 4. The upper estimates follow immediately from Lemma 9.4 in [3]. Only the lower estimate in (9) was not proved in [3]. Briefly, the lower estimate follows from the boundedness of F . At the neck, one has

$$K = \frac{\psi_{ss}}{\psi}, \quad L = \frac{1}{\psi^2}, \quad \psi = (1 + o(1))\sqrt{2(n-1)(T-t)},$$

so that $(K/L)\log L \leq C$ implies

$$0 \leq \psi_{ss} \leq \frac{C}{\tau\psi}.$$

The radius $\psi(0, t)$ of the neck satisfies

$$\psi_t = \psi_{ss} - \frac{n-1}{\psi} \leq -\frac{n-1-C(\tau^{-1})}{\psi},$$

which upon integration yields

$$\psi(0, t) \geq \left(1 - \frac{\tilde{C}}{\tau}\right) \sqrt{2(n-1)(T-t)}$$

where $\tilde{C}$ only depends on F^* and n . This implies the lower estimates in the Lemma. $\square$

2.3. Transition to commuting variables. On most of the neck, the quantity $v = o(1)$. So it is natural to drop the quadratic terms in (7), resulting in the deceptively simple equation

$$v_\tau = v_{\sigma\sigma} + v$$

for v . Here the partial derivatives ∂_τ and ∂_σ do not commute. To work with commuting variables, we regard v as function of σ . In other words, we define

$$(15) \quad \sigma(x, t) = \frac{1}{\sqrt{T-t}} \int_0^x ds = e^{-\tau/2} \int_0^x ds,$$

and consider a function $\tilde{v}(\sigma, \tau)$ such that $\tilde{v}(\sigma(x, \tau), \tau) \equiv v(x, \tau)$. We shall abuse notation and write v for both quantities v and $\tilde{v}$. This is only ambiguous when we take time derivatives, in which case we write

$$\left. \frac{\partial v}{\partial \tau} \right|_{\sigma \text{ const}} = \frac{\partial \tilde{v}}{\partial \tau} \quad \text{and} \quad \left. \frac{\partial v}{\partial \tau} \right|_{x \text{ const}} = \frac{\partial v}{\partial \tau}.$$

The two are related by

$$\left. \frac{\partial v}{\partial \tau} \right|_{\sigma \text{ const}} = \left. \frac{\partial v}{\partial \tau} \right|_{x \text{ const}} - \frac{\partial \sigma}{\partial \tau} \frac{\partial v}{\partial \sigma}.$$

Lemma 5. *Assuming reflection symmetry of the metric, i.e. $\psi(-s, t) \equiv \psi(s, t)$, one has*

$$\frac{\partial \sigma}{\partial \tau} = \frac{\sigma}{2} + nJ(\sigma, \tau),$$

where

$$J(\sigma, \tau) = \int_0^\sigma \frac{v_{\sigma\sigma}}{1+v} d\sigma = \frac{v_\sigma}{1+v} + \int_0^\sigma \frac{v_\sigma^2}{(1+v)^2} d\sigma.$$

Proof. By definition one has $\sigma(x, \tau) = e^{\tau/2} s(x, T - e^{-\tau})$. Hence

$$\sigma_\tau = \frac{1}{2}\sigma + e^{\tau/2} s_t(x, T - e^{-\tau}) e^{-\tau} = \frac{1}{2}\sigma + e^{-\tau/2} s_t(x, T - e^{-\tau}).$$

One also has [3, §10]

$$\frac{\partial s}{\partial t} = \int_0^s \frac{n\psi_{ss}}{\psi} ds = n \left\{ \frac{\psi_s}{\psi} + \int_0^s \frac{\psi_s^2}{\psi^2} ds \right\}.$$

Using the relations (2) and (3) between $u = 1 + v$ and ψ , and s and σ , one finds the stated expression for $\partial_\tau \sigma$. $\square$

From here on we will consider v as a function of σ and τ . All τ derivatives are intended to be time derivatives with σ kept constant. It follows from (7) that v satisfies

$$(16) \quad v_\tau = v_{\sigma\sigma} - \frac{\sigma}{2} v_\sigma + v - nJ(\sigma, \tau) v_\sigma + \frac{(n-1)v_\sigma^2 - \frac{1}{2}v^2}{1+v}.$$

We write this equation as

$$v_\tau = A(v) + N(v),$$

where A is the linear differential operator

$$A = \left(\frac{\partial}{\partial \sigma} \right)^2 - \frac{\sigma}{2} \frac{\partial}{\partial \sigma} + 1,$$

and $N(v)$ represents the remaining (nonlinear) terms in (16),

$$N(v) = -nv_\sigma \int_0^\sigma \frac{v_{\sigma\sigma}}{1+v} d\sigma + \frac{(n-1)v_\sigma^2 - \frac{1}{2}v^2}{1+v}.$$

Integrate by parts to get

$$(17) \quad N(v) = -nv_\sigma \int_0^\sigma \frac{v_\sigma^2}{(1+v)^2} d\sigma - \frac{v_\sigma^2 + \frac{1}{2}v^2}{1+v}.$$

Lemma 6. *For all $|\sigma| \leq e^{\delta\tau}$, one has*

$$|N(v)| \leq C \frac{1 + \sigma^4}{\tau^2}.$$

Proof. This follows from Lemma 4. In fact, Lemma 4 implies that

$$(18) \quad |v| \leq C \frac{1 + \sigma^2}{\tau} \text{ and } |v_\sigma| \leq C \frac{1 + |\sigma|}{\tau}$$

for $|\sigma| \leq 2\sqrt{\tau}$. For larger σ , one finds other estimates for v and v_σ , which are stronger than (18). Using (18) for all $|\sigma| \leq e^{\delta\tau}$, one arrives at our estimate for $N(v)$. $\square$

2.4. The linearized equation. If we ignore the nonlinear terms in (16), then we see that the small quantity v satisfies $v_\tau = A(v)$. Even though v is not defined for all $s \in \mathbb{R}$, we know from our estimate on the neck that v is defined for $|\sigma| \leq e^{\delta\tau}$, for some $\delta = \delta(F^*, n)$. In the first analysis, we assume that v is in fact a solution of the Cauchy problem $v_\tau = A(v)$. The operator A is self-adjoint in the Hilbert space

$$\mathcal{H} = \left\{ v \in L^2 \left(\mathbb{R}, e^{-\sigma^2/4} d\sigma \right) \mid v(\sigma) \equiv v(-\sigma) \right\}.$$

It has pure point spectrum, with eigenvalues $\lambda_m = 1 - m$ for $m = 0, 1, 2, \dots$. The eigenfunction corresponding to the eigenvalue λ_m is an even *Hermite polynomial* of degree $2m$. Normalizing the eigenfunction so that the coefficient of its highest order term is 1, we set

$$h_{2m}(\sigma) = \sigma^{2m} - \frac{(2m)!}{1!(2m-2)!} \sigma^{2m-2} + \frac{(2m)!}{2!(2m-4)!} \sigma^{2m-4} - \dots$$

In particular,

$$(19) \quad h_0(\sigma) = 1, \quad h_2(\sigma) = \sigma^2 - 2, \quad h_4(\sigma) = \sigma^4 - 12\sigma^2 + 12.$$

2.5. Eigenfunction decomposition of v . Let $\eta \in C^\infty(\mathbb{R})$ be an even bump function with $\eta(z) = 0$ for $z \geq 1\frac{1}{4}$ and $\eta(z) = 1$ for $z \leq 1$. We then define

$$(20) \quad V(\sigma, \tau) = \eta(\sigma e^{-\delta\tau/2}) v(\sigma, \tau).$$

For $|\sigma| \geq \frac{5}{4}e^{\delta\tau/2}$, we set $V \equiv 0$. Then $V(\cdot, \tau) \in \mathcal{H}$, and $v \equiv V$ for $|\sigma| \leq e^{\delta\tau/2}$.

A computation shows that

$$(21) \quad V_\tau - A(V) = \eta N(v) + E,$$

where the “error term” is

$$E = \left(\eta_\tau - \eta_{\sigma\sigma} + \frac{\sigma}{2} \eta_\sigma \right) v - 2\eta_\sigma v_\sigma.$$

Lemma 7. *The error term E vanishes except when $e^{\delta\tau/2} \leq |\sigma| \leq \frac{5}{4}e^{\delta\tau/2}$. When $E \neq 0$, one has*

$$|E(\sigma, \tau)| \leq C|\sigma|,$$

where the constant C only depends on F^* and n . One also has

$$\|E(\cdot, \tau)\| \leq C \exp\left(-e^{\delta\tau/2}\right),$$

where the constant C again only depends on F^* and n .

Proof. The pointwise estimate follows from the boundedness of η_τ , $\sigma\eta_\sigma$, and $\eta_{\sigma\sigma}$, as well as the pointwise bounds for v provided by Lemma 4. Given the pointwise bounds for E , one finds that

$$\|E\|^2 \leq C^2 \int_{e^{\delta\tau/2}}^{\infty} \sigma^2 e^{-\sigma^2/4} d\sigma,$$

from which the $\mathcal{H}$ norm estimate follows. $\square$

For each τ , the function $V(\cdot, \tau)$ belongs to $\mathcal{H}$. So we can consider the splitting into mutually orthogonal terms determined by

$$(22) \quad V(\sigma, \tau) = a_0(\tau)h_0(\sigma) + a_2(\tau)h_2(\sigma) + W(\sigma, \tau).$$

Our pointwise estimate (14) for v suggests that for large τ these terms will decay like τ^{-1} . This turns out to be the case for the middle term $a_2(\tau)h_2(\tau)$, but the other two terms are in fact smaller. We will show in the next few paragraphs that they decay like τ^{-2} . This establishes $a_2(\tau)h_2(\tau)$ as the dominant term for large τ .

2.6. Easy estimates for a_0 and a_2 . Lemma 1 and in particular (14) imply that

$$|V(\sigma, \tau)| \leq \frac{C}{\tau}(1 + \sigma^2)$$

for all $\sigma \in \mathbb{R}$, and all $\tau \geq \tau_0$. Upon taking the $\mathcal{H}$ -inner product with h_0 and h_2 this gives us

$$(23) \quad |a_0(\tau)| + |a_2(\tau)| \leq \frac{C}{\tau}$$

for all $\tau \geq \tau_0$.

2.7. Decay of a_0 . Since $\|h_0\|^2 a_0(\tau) = (h_0, V)_\mathcal{H}$, we have

$$\begin{aligned} \|h_0\|^2 a_0'(\tau) &= (h_0, V_\tau)_\mathcal{H} \\ &= (h_0, A(V) + \eta N(v) + E)_\mathcal{H} \\ &= (h_0, V)_\mathcal{H} + (h_0, \eta N(v) + E)_\mathcal{H} \\ &= \|h_0\|^2 a_0(\tau) + (h_0, \eta N(v) + E)_\mathcal{H}. \end{aligned}$$

Hence

$$a_0'(\tau) - a_0(\tau) = \left(\frac{h_0}{\|h_0\|^2}, \eta N(v) + E \right)_\mathcal{H}.$$

Using our estimates for $N(v)$ and E , we conclude that

$$f_0(\tau) \doteq a_0'(\tau) - a_0(\tau)$$

satisfies

$$|f_0(\tau)| \leq \frac{C(F^*, n)}{\tau^2}.$$

The variation of constants formula tells us that

$$a_0(\tau) = - \int_\tau^\infty e^{\tau-\tau'} f_0(\tau') d\tau',$$

and thus that

$$|a_0(\tau)| \leq \frac{C(F^*, n)}{\tau^2}.$$

2.8. Decay of W . Since W lies in the stable space of the operator A , i.e. the space spanned by those eigenfunctions with negative eigenvalues, we expect W to decay according to the slowest stable eigenvalue (i.e. $\sim e^{-\tau}$), or else to decay like the “forcing term” $\eta N(v) + E$. In this situation, the forcing term is dominant, and we have the following.

Lemma 8. *For any given $\tau_0 > 0$, one has for all $\tau > \tau_0$*

$$\|W(\cdot, \tau)\| \leq e^{\tau_0 - \tau} \|W(\cdot, \tau_0)\| + \frac{C}{\tau^2}.$$

Consequently, one also has

$$\|W(\cdot, \tau)\| \leq \frac{C}{\tau^2} (1 + W_0).$$

Here the constant C only depends on F^ and n , and*

$$W_0 \doteq e^{\tau_0} \|W(\cdot, \tau_0)\|.$$

Proof. Since $\|W\|^2 = (W, W) = (W, V)$ and $(W, A(V))_{\mathcal{H}} = (W, A(W))_{\mathcal{H}}$, one has

$$\begin{aligned} \|W\| \frac{d\|W\|}{d\tau} &= \frac{1}{2} \frac{d}{d\tau} \|W\|^2 \\ &= \frac{1}{2} \frac{d}{d\tau} (W, V)_{\mathcal{H}} \\ &= (W, V_{\tau})_{\mathcal{H}} \\ &= (W, A(V) + \eta N(v) + E)_{\mathcal{H}} \\ &= (W, A(W))_{\mathcal{H}} + (W, \eta N(v) + E)_{\mathcal{H}} \\ &\leq -\|W\|^2 + \|W\| \cdot \|\eta N(v) + E\|, \end{aligned}$$

whence

$$(24) \quad \frac{d\|W\|}{d\tau} \leq -\|W\| + \|\eta N(v) + E\|.$$

From Lemma 6 we conclude that $|\eta N(v)| \leq C(1 + \sigma^4)\tau^{-2}$ for all σ , and hence that

$$\|\eta N(v)\| \leq \frac{C}{\tau^2}.$$

For E , we use Lemma 7 together with the calculus inequality $Ce^{-e^{\delta\tau/2}} \leq C'\tau^{-2}$. We apply all this to (24) and conclude that

$$\frac{d\|W\|}{d\tau} \leq -\|W\| + \frac{C}{\tau^2}.$$

Therefore

$$\frac{d(e^{\tau}\|W\|)}{d\tau} \leq Ce^{\tau}\tau^{-2}.$$

Integration leads to

$$e^{\tau}\|W(\cdot, \tau)\| \leq e^{\tau_0}\|W(\cdot, \tau_0)\| + C \int_{\tau_0}^{\tau} \frac{e^{\tilde{\tau}}}{\tilde{\tau}^2} d\tilde{\tau} \leq e^{\tau_0}\|W(\cdot, \tau_0)\| + \frac{\tilde{C}}{\tau^2} e^{\tau},$$

as claimed. The second estimate for $\|W\|$ follows directly from $\sup \tau^2 e^{-\tau} < \infty$. $\square$

2.9. Derivative estimates for W . We use the regularizing effect of the heat equation to bootstrap the estimates of Lemma 8 by one space derivative.

Lemma 9. *Let $\tau_0 > 0$ be as in Lemma 8. Then for all $\tau > \tau_0 + 1$, one has*

$$\|W_\sigma(\cdot, \tau)\| \leq C(1 + W_0)\tau^{-2},$$

where the constant C only depends on F^* and n , and as before, $W_0 = e^{\tau_0}\|W(\cdot, \tau_0)\|$.

Proof. Note that

$$Au = u_{\sigma\sigma} - \frac{1}{2}\sigma u_\sigma + u = e^{\sigma^2/4}(e^{-\sigma^2/4}u_\sigma)_\sigma$$

so that for $u \in D(A)$ one has

$$-(u, Au)_\mathcal{H} = \int_{\mathbb{R}} e^{-\sigma^2/4} \{u_\sigma^2 - u^2\} d\sigma.$$

Thus

$$(25) \quad \|\partial_\sigma u\|^2 = \|u\|^2 - (u, Au).$$

If $u \perp h_0, h_2$ then $(u, -Au)_\mathcal{H} \geq 0$, and we have

$$\|\partial_\sigma u\| \leq \|u\| + \sqrt{(u, -Au)_\mathcal{H}}.$$

Applying the spectral theorem to the selfadjoint operator A on the Hilbert space $\{h_0, h_2\}^\perp$ in order to estimate $(u, Ae^{\theta A}u)$, one concludes that

$$(26) \quad \|\partial_\sigma e^{\theta A}u\| \leq \frac{C}{\sqrt{\theta}}\|u\|$$

holds for $0 < \theta \leq 1$, and all $u \in D(A)$.

The stable component W of V satisfies

$$W_\tau = A(W) + f(\tau), \text{ where } f(\tau) \doteq \mathcal{P}(\eta N(v) + E),$$

and where $\mathcal{P}$ is the $\mathcal{H}$ -orthogonal projection onto the space $\{u \in \mathcal{H} \mid u \perp h_0, h_2\}$. By the variation of constants formula one then has, for all $\tau \geq \tau_0 + 1$

$$W(\tau) = e^A W(\tau - 1) + \int_0^1 e^{\theta A} f(\tau - \theta) d\theta.$$

Since $f(\tau) \perp \{h_0, h_2\}$, we can use (26) to get

$$\begin{aligned} \|\partial_\sigma W(\tau)\| &\leq \|\partial_\sigma e^A W(\tau - 1)\| + \int_0^1 \|\partial_\sigma e^{\theta A} f(\tau - \theta)\| d\theta \\ &\leq C\|W(\tau - 1)\| + \int_0^1 \frac{C}{\sqrt{\theta}} \|f(\tau - \theta)\| d\theta. \end{aligned}$$

The combination of Lemmas 6 and 7 again gives $\|f(\tau)\| \leq C\tau^{-2}$. The integral can therefore be bounded by $C\tau^{-2}$. Combining this with our previous estimate for $\|W(\tau - 1)\|$ from Lemma 8, we get the inequality in Lemma 9. $\square$

2.10. A bound for $\|\sigma W\|$. If $\varphi \in C_c^1(\mathbb{R})$, then one has the identity

$$\int_{\mathbb{R}} e^{-\sigma^2/4} 2\varphi(\sigma) \sigma \varphi'(\sigma) d\sigma = \int_{\mathbb{R}} e^{-\sigma^2/4} (-2\varphi(\sigma)^2 + \frac{1}{2}\sigma^2 \varphi(\sigma)^2) d\sigma$$

which after rearranging leads to

$$\begin{aligned} \|\sigma\varphi\|^2 &= 4 \int_{\mathbb{R}} e^{-\sigma^2/4} (\sigma\varphi\varphi' + \varphi^2) d\sigma \\ &= 4 \{(\sigma\varphi, \varphi')_{\mathcal{H}} + \|\varphi\|^2\} \\ &\leq \frac{1}{2}\|\sigma\varphi\|^2 + 8\|\varphi'\|^2 + 4\|\varphi\|^2 \\ &\leq 16 \{\|\varphi'\|^2 + \|\varphi\|^2\}. \end{aligned}$$

An approximation argument leads to

Lemma 10. *For any $\varphi \in \mathcal{H}$ with $\varphi_\sigma \in \mathcal{H}$ one has $\sigma\varphi \in \mathcal{H}$, while*

$$\|\sigma\varphi(\sigma)\| \leq 4 \{\|\varphi_\sigma\| + \|\varphi\|\}.$$

In particular, W satisfies

$$(27) \quad \|\sigma W\| \leq C(1 + W_0)\tau^{-2}$$

for all $\tau \geq \tau_0 + 1$.

2.11. Pointwise estimates for W . A Sobolev inequality says that the bound on $\|\partial_\sigma W\|$ implies a pointwise bound for W . Due to the exponential weight in the norm this bound is not uniform. One has

Lemma 11. *Any $\varphi \in \mathcal{H}$ with $\partial_\sigma \varphi \in \mathcal{H}$ is a continuous function, which satisfies*

$$e^{-\sigma^2/8} |\varphi(\sigma)| \leq C(\|\varphi\| + \|\partial_\sigma \varphi\|).$$

Hence W satisfies

$$|W(\sigma, \tau)| \leq \frac{C(1 + W_0)}{\tau^2} e^{\sigma^2/8}.$$

Proof. Assuming $\varphi \in C_c^1(\mathbb{R})$ one has

$$\begin{aligned} e^{-\sigma^2/4} \varphi(\sigma)^2 &= \int_{-\infty}^{\sigma} e^{-\sigma^2/4} \left[-\frac{\sigma}{2} \varphi^2 + 2\varphi\varphi_\sigma \right] d\sigma \\ &\leq \|\sigma\varphi\| \|\varphi\| + \|\varphi\| \|\varphi_\sigma\| \\ &\leq C \{\|\varphi\|^2 + \|\varphi_\sigma\|^2\}. \end{aligned}$$

So the pointwise estimate holds for $\varphi \in C_c^1$. By approximation it also holds for all $\varphi \in \mathcal{H}$ with $\partial_\sigma \varphi \in \mathcal{H}$. $\square$

Lemma 11 says that W decays pointwise like τ^{-2} , and uniformly on bounded σ intervals. The following pointwise bound for W only gives a τ^{-1} decay rate, but it is stronger for large values of σ . The estimate also does not depend on W_0 .

Lemma 12. *For some constant $C = C(F^*, n) < \infty$ one has for all $\tau > \tau_0 + 1$*

$$|W(\sigma, \tau)| \leq \frac{C}{\tau} (1 + \sigma^2).$$

Proof. We have, by definition, $W(\sigma, \tau) = V(\sigma, \tau) - a_0(\tau)h_0(\sigma) - a_2(\tau)h_2(\sigma)$. Using our pointwise bounds for V , a_0 , a_2 , and $h_0(\sigma) \equiv 1$, $h_2(\sigma) \equiv \sigma^2 - 2$, one quickly gets the stated estimate for W . $\square$

2.12. Pointwise estimate for W_σ . By definition we have $W(\tau, \sigma) = V(\tau, \sigma) - a_0(\tau)h_0(\sigma) - a_2(\tau)h_2(\sigma)$. Differentiate and keep in mind that $h'_0(\sigma) = 0$ and $h'_2(\sigma) = 2\sigma$. We get $W_\sigma = V_\sigma - 2a_2(\tau)\sigma$. Using the pointwise derivative bounds (11) and (12) for u , one then gets

$$(28) \quad |W_\sigma| \leq \frac{C}{\tau}(1 + |\sigma|)$$

which holds for all $\tau \geq \tau_0$ and $\sigma \in \mathbb{R}$.

2.13. An equation for a_2 . As with a_0 , we have

$$\begin{aligned} \|h_2\|^2 a'_2(\tau) &= (h_2, V_\tau)_\mathcal{H} \\ &= (h_2, A(V) + \eta N(v) + E)_\mathcal{H} \\ &= (A(h_2), V)_\mathcal{H} + (h_2, \eta N(v) + E)_\mathcal{H} \\ &= (h_2, \eta N(v) + E)_\mathcal{H}. \end{aligned}$$

We use (17) and the identity $1/(1+v) = 1 - v/(1+v)$ to rewrite $N(v)$ as

$$(29) \quad N(v) = N_2(v) + N_3(v), \text{ where } \begin{cases} N_2(v) = -(v_\sigma^2 + \frac{1}{2}v^2) \\ N_3(v) = -nv_\sigma \int_0^\sigma \frac{v_\sigma^2}{(1+v)^2} d\sigma + \frac{vv_\sigma^2 + \frac{1}{2}v^3}{1+v}. \end{cases}$$

Here, $N_2(v)$ is the purely quadratic part, while $N_3(v)$ contains the cubic and higher order terms.

Lemma 13. *There is a constant $C = C(F^*, n)$ such that for all $\tau \geq \tau_0$*

$$\|\eta N_3(v)\| \leq \frac{C}{\tau^3}.$$

Proof. The pointwise estimates for v and v_σ from (14) imply that

$$|v| \leq C \frac{1 + \sigma^2}{\tau}, \quad |v_\sigma| \leq C \frac{1 + |\sigma|}{\tau}$$

whenever $|\sigma| \leq e^{\delta\tau}$. Hence we have

$$|N_3(v)| \leq C \frac{1 + \sigma^6}{\tau^3} \text{ for } |\sigma| \leq e^{\delta\tau}.$$

(The term with v^3 contributes the highest power in σ .)

This implies the Lemma. $\square$

We continue with our computation of $a'_2(\tau)$. We have

$$\begin{aligned} \|h_2\|^2 a'_2(\tau) &= (h_2, \eta N(v) + E)_\mathcal{H} \\ &= (h_2, \eta N_2(v) + \eta N_3(v) + E)_\mathcal{H} \\ &= (h_2, \eta N_2(v))_\mathcal{H} + (h_2, \eta N_3(v) + E)_\mathcal{H}. \end{aligned}$$

The last term satisfies

$$|(h_2, \eta N_3(v) + E)_{\mathcal{H}}| \leq \|h_2\| (\|\eta N_3(v)\| + \|E\|) \leq \frac{C}{\tau^3}.$$

We would now like to replace the v in the quadratic expression $N_2(v)$ by V , and then by the dominant term in (22), i.e. $a_2(\tau)h_2(\sigma)$. Before we do this, we estimate the errors produced by these replacements.

Lemma 14. *There is a constant $C = C(F^*, n)$ such that for all $\tau \geq \tau_0$, one has*

$$\|\eta(N_2(v) - N_2(V))\| \leq Ce^{-e^{\delta\tau/2}}.$$

In other words $\eta(N_2(v) - N_2(V))$ satisfies the same super-exponentially small estimate as the error term E . (See Lemma 7.)

Proof. Since $V = \eta \cdot v$, a direct calculation gives

$$\eta(N_2(v) - N_2(V)) = \eta \left\{ (\eta v)_\sigma^2 - v_\sigma^2 \right\} + \frac{1}{2}\eta(\eta^2 - 1)v^2.$$

Using the pointwise estimates for v and v_σ with the fact that $\eta(N_2(v) - N_2(V))$ is supported in the region $e^{\delta\tau/2} \leq |\sigma| \leq \frac{5}{4}e^{\delta\tau/2}$, one arrives at the stated estimate for the $\mathcal{H}$ norm of $\eta(N_2(v) - N_2(V))$. $\square$

We therefore have

$$\|h_2\|_{\mathcal{H}}^2 a'_2(\tau) = (h_2, \eta N_2(V))_{\mathcal{H}} + (h_2, \eta(N_2(v) - N_2(V)))_{\mathcal{H}} + (h_2, \eta N_3(v) + E)_{\mathcal{H}}.$$

If we write $N_2[v, w] = -v_\sigma w_\sigma - \frac{1}{2}vw$ for the natural symmetric bilinear expression with $N_2(v) = N_2[v, v]$, then the first term above can be written as

$$\begin{aligned} (h_2, \eta N_2(V))_{\mathcal{H}} &= (\eta h_2, N_2(a_2 h_2 + a_0 h_0 + W))_{\mathcal{H}} \\ &= (a_2)^2 (\eta h_2, N_2(h_2))_{\mathcal{H}} \\ &\quad + 2a_2 (\eta h_2, N_2[h_2, a_0 h_0 + W])_{\mathcal{H}} \\ &\quad + (\eta h_2, N_2(a_0 h_0 + W))_{\mathcal{H}}, \end{aligned}$$

which is a quadratic polynomial in a_2 . Adding in the omitted error terms, we reach the following observation.

Lemma 15.

$$a'_2(\tau) = K(\tau)a_2(\tau)^2 + 2L(\tau)a_2(\tau) + M(\tau),$$

in which

$$K(\tau) = \frac{1}{\|h_2\|^2} (\eta h_2, N_2(h_2))_{\mathcal{H}},$$

$$L(\tau) = \frac{1}{\|h_2\|^2} (\eta h_2, N_2[h_2, a_0 h_0 + W])_{\mathcal{H}},$$

$$M(\tau) = \frac{1}{\|h_2\|^2} \{ (\eta h_2, N_2(a_0 h_0 + W))_{\mathcal{H}} + (h_2, \eta(N_2(v) - N_2(V)))_{\mathcal{H}} + (h_2, \eta N_3(v) + E)_{\mathcal{H}} \}.$$

Lemma 16.

$$K(\tau) = -8 + \mathcal{O}\left(e^{-e^{\delta\tau/2}}\right).$$

Proof. Since $N_2(v) = -v_\sigma^2 - \frac{1}{2}v^2$, the explicit expressions (19) for h_0 , h_2 , and h_4 give us

$$N_2(h_2) = -\frac{1}{2}\sigma^4 - 2\sigma^2 - 2 = -\frac{1}{2}h_4 - 8h_2 - 12h_0.$$

Hence $(h_2, N_2(h_2))_{\mathcal{H}} = -8\|h_2\|^2$. This implies

$$K(\tau) = \frac{(h_2, N_2(h_2))_{\mathcal{H}}}{\|h_2\|^2} + \frac{((1-\eta)h_2, N_2(h_2))_{\mathcal{H}}}{\|h_2\|^2}.$$

The first term is -8 . The other term can be written as the integral of a function with fixed polynomial growth, which is supported in the region $|\sigma| \geq e^{\delta\tau/2}$. This leads quickly to the stated estimate. $\square$

Lemma 17. $L(\tau) = \mathcal{O}(\tau^{-2})$, i.e. there is a constant $C = C(F^*, n)$ such that for all $\tau \geq \tau_0 + 1$ one has $|\tau^2 L(\tau)| \leq C$.

Proof. Using (bi)linearity of $N_2[v, \tilde{v}]$, we get

$$\begin{aligned} \|h_2\|^2 L(\tau) &= (\eta h_2, N_2[h_2, a_0 h_0 + W])_{\mathcal{H}} \\ &= a_0(\tau) (h_2, \eta N_2(h_2, h_0))_{\mathcal{H}} + (h_2, \eta N_2(h_2, W))_{\mathcal{H}}. \end{aligned}$$

The first term is $\mathcal{O}(\tau^{-2})$. For the other term, we have

$$\begin{aligned} |(h_2, \eta N_2(h_2, W))_{\mathcal{H}}| &= |(\eta h_2, h'_2(\sigma)W_\sigma + \frac{1}{2}h_2 W)_{\mathcal{H}}| \\ &\leq \|\eta h_2 h'_2\| \|W_\sigma\| + C\|W\| \\ &\leq \frac{C}{\tau^2} \end{aligned}$$

by our estimates for $\|W\|$ and $\|\partial_\sigma W\|$. $\square$

Lemma 18.

$$M(\tau) \leq C \left\{ \frac{e^{\tau_0 - \tau}}{\tau} \|W(\cdot, \tau_0)\| + \frac{1}{\tau^3} \right\}$$

In fact one can show that $M(\tau) \leq C_\alpha \tau^{-(4-\alpha)}$ for any $\alpha > 0$, but we will only need the estimate with $\alpha = 1$.

Proof. In Lemma 15, $\|h_2\|^2 M(\tau)$ is defined as the sum of three terms. We now estimate them one by one.

The middle term is

$$|(h_2, \eta(N_2(v) - N_2(V)))_{\mathcal{H}}| \leq \|h_2\| \|\eta(N_2(v) - N_2(V))\| \leq C e^{-e^{\delta\tau/2}},$$

by Lemma 14.

The third term is bounded by

$$|(h_2, \eta N_3(v) + E)_{\mathcal{H}}| \leq C (\|\eta N_3(v)\| + \|E\|) \leq \frac{C}{\tau^3},$$

because of Lemma 13 and the super-exponential estimate for E in Lemma 7.

The first term can itself be split into three terms:

$$(30) \quad (\eta h_2, N_2(a_0 h_2 + W))_{\mathcal{H}} = a_0(\tau)^2 (\eta h_2, N_2(h_2))_{\mathcal{H}} + 2a_0(\tau) (\eta h_2, N_2[h_2, W])_{\mathcal{H}} + (\eta h_2, N_2(W))_{\mathcal{H}}.$$

The first of these terms is $\mathcal{O}(\tau^{-4})$, since $a_0(\tau) = \mathcal{O}(\tau^{-2})$.

From

$$\begin{aligned}
(\eta h_2, N_2[h_2, W])_{\mathcal{H}} &= (\eta h_2, -h_2'(\sigma)W_\sigma - \tfrac{1}{2}h_2W)_{\mathcal{H}} \\
&= -(\eta h_2 h_2', W_\sigma)_{\mathcal{H}} - \tfrac{1}{2}(\eta h_2(\sigma)^2, W)_{\mathcal{H}} \\
&= \mathcal{O}(\|W_\sigma\| + \|W\|) \\
&= \mathcal{O}(\tau^{-2})
\end{aligned}$$

and $a_0(\tau) = \mathcal{O}(\tau^{-2})$, we deduce that the middle term in (30) is also $\mathcal{O}(\tau^{-4})$.

We are left with the last term in (30). This term can be expanded as

$$(31) \quad (\eta h_2, N_2(W))_{\mathcal{H}} = -(\eta h_2, W_\sigma^2)_{\mathcal{H}} - (\eta h_2, \tfrac{1}{2}W^2)_{\mathcal{H}}.$$

The second term satisfies

$$\begin{aligned}
\left| \int_{\mathbb{R}} e^{-\sigma^2/4} \eta h_2(\sigma) W(\tau, \sigma)^2 d\sigma \right| &\leq C(\|W\|^2 + \|\sigma W\|^2) \\
&\leq C \left\{ e^{2(\tau_0 - \tau)} \|W(\cdot, \tau_0)\|^2 + \frac{1}{\tau^4} \right\},
\end{aligned}$$

by Lemmas 8 and 10.

We bound the first term in (31) as follows:

$$\begin{aligned}
|(\eta h_2, W_\sigma^2)_{\mathcal{H}}| &= \left| \int_{\mathbb{R}} e^{-\sigma^2/4} \eta h_2(\sigma) W_\sigma^2 d\sigma \right| \\
&\leq C \left| \int_{\mathbb{R}} e^{-\sigma^2/4} (1 + \sigma^2) W_\sigma^2 d\sigma \right| \\
&\leq C \|(1 + \sigma)^2 W_\sigma\| \|W_\sigma\|.
\end{aligned}$$

The pointwise estimate (28) bounds the first factor by

$$\|(1 + \sigma)^2 W_\sigma\| \leq \frac{C}{\tau},$$

while Lemma 9 implies

$$\|W_\sigma\| \leq C \left\{ e^{\tau_0 - \tau} \|W(\cdot, \tau_0)\| + \frac{1}{\tau^2} \right\}.$$

Combined, these last two estimates imply Lemma 18. $\square$

2.14. Dichotomy for the decay of $a_2(\tau)$. We have shown that

$$(32) \quad \frac{da_2(\tau)}{d\tau} = -8a_2(\tau)^2 + M^*(\tau),$$

where

$$|M^*(\tau)| \leq C(1 + W_0)\tau^{-3}.$$

Hence

$$\tau \frac{d(\tau a_2(\tau))}{d\tau} = (\tau a_2(\tau)) - 8(\tau a_2(\tau))^2 + \tau^2 M^*(\tau).$$

Lemma 19. *The coefficient $a_2(\tau)$ satisfies either*

$$(33) \quad a_2(\tau) \leq C(1 + W_0)\tau^{-2}$$

for all $\tau \geq \tau_0 + 1$, or else

$$(34) \quad \lim_{\tau \rightarrow \infty} \tau a_2(\tau) = \frac{1}{8}.$$

2.15. Pinching time. We will show below that alternative (34) does indeed occur, namely that there exists an open set of solutions matching the asymptotic profile we have constructed. Our first step is to improve the estimate obtained in Lemma 6.1 of [3]. Let $r(t) \equiv \psi(0, t)$ denote the radius of the neck, and recall that

$$K = \frac{\psi_{ss}}{\psi} \quad \text{and} \quad L = \frac{1 - \psi_s^2}{\psi^2}$$

denote the sectional curvatures. Define $\lambda = 2 - \log L_{\min}(0)$, so that we may write the cylindricity F as

$$F = \frac{K}{L}(\lambda + \log L) = \frac{\psi\psi_{ss}}{1 - \psi_s^2} \left(\lambda + \log \frac{1 - \psi_s^2}{\psi^2} \right).$$

Recall that there exists $F^* \geq n - 1$ depending only on $g(0)$ such that $0 \leq F \leq F^*$ on the neck. Writing the evolution equation for the radius

$$\frac{dr}{dt} = \left[\psi_{ss} - (n - 1) \frac{1 - \psi_s^2}{\psi} \right]_{s=0} = \psi_{ss} - \frac{n - 1}{r}$$

in the form

$$(35) \quad \frac{d}{dt}(r^2) = -2(n - 1) + 2 \frac{F}{\lambda + \log \frac{1}{r^2}}$$

then proves that

$$(36) \quad -2(n - 1) \leq \frac{d}{dt}(r^2) \leq -2(n - 1) + 2 \frac{F^*}{\lambda + \log \frac{1}{r^2}}.$$

This observation lets us prove that the singularity time depends continuously on the initial conditions.

Lemma 20. *For rotationally symmetric metrics $g(t)$ that develop a neckpinch, the pinching time T is a continuous function of the initial metric g_0 .*

Proof. Given a metric g_0 , let $T_0 = T(g_0)$ denote the singularity time of the solution $g(t)$ with initial data g_0 .

To prove lower semicontinuity, let $T_- < T_0$ be given. Then $\{g(t) : 0 \leq t \leq T_-\}$ is a smooth solution of Ricci flow. Because regular solutions of parabolic equations depend continuously on their initial data, a solution $\tilde{g}(t)$ will remain smooth for $0 \leq t \leq T_-$, provided that its initial data $\tilde{g}_0$ is sufficiently close to g_0 . This implies that the singularity time $\tilde{T}$ of $\tilde{g}(\cdot)$ satisfies $\tilde{T} > T_-$.

To prove upper semicontinuity, let $T_+ > T_0$ be given. At any time $T_* < T_0$, observe that for all $\tilde{g}_0$ sufficiently close to g_0 , the radius $\tilde{r}$ of the solution $\tilde{g}(t)$ satisfies $\tilde{r}(T_*)^2 \leq 2r(T_*)^2$, and its cylindricity $\tilde{F}$ satisfies $\tilde{F} \leq \tilde{F}^* \leq 2F^*$. By

taking T_* close enough to T_0 , one may by (36) ensure that any solution $\tilde{g}(t)$ with initial data $\tilde{g}_0$ satisfies

$$\frac{d}{dt}(\tilde{r}^2) \leq -2(n-1) + \frac{4F^*}{\lambda + \log \frac{1}{2r(T_*)^2}} \leq -(n-1)$$

for all $t \in (T_*, \tilde{T})$, where $\tilde{T}$ is the pinching time of $\tilde{g}(t)$. So for $t \in (T_*, \tilde{T})$, it follows that

$$\begin{aligned} \tilde{r}(t)^2 &\leq \tilde{r}(T_*)^2 - (n-1)(t - T_*) \\ &\leq 2r(T_*)^2 - (n-1)(t - T_*) \\ &\leq 4(n-1)(T_0 - T_*) - (n-1)(t - T_*). \end{aligned}$$

Here we used the fact that $r(t)^2 \leq 2(n-1)(T_0 - t)$. Choose $T_* < T_0$ closer to T_0 if necessary so that $T_* + 4(T_0 - T_*) < T_+$. Then if $t \geq T_* + 4(T_0 - T_*)$, one has $\tilde{r}(t)^2 \leq 0$. This implies that $\tilde{T} \leq T_* + 4(T_0 - T_*) < T_+$. $\square$

2.16. Initial data. Given a target time T , we now construct initial data $u = 1 + v$ at $t = t_0 < T$ that will become singular at T and satisfy the asymptotic profile (34). In the next subsection, we shall obtain appropriate estimates for the function $V(\sigma, \tau) = \eta(e^{-\delta\tau/2}\sigma) \cdot v(\sigma, \tau) = \sum_{k=0}^{\infty} a_{2k}(\tau)h_{2k}(\sigma)$ corresponding to this construction.

Let $\varepsilon \in (0, \frac{1}{4\sqrt{n-1}})$ be a constant to be chosen later. Near the developing neck-pinch, we impose the initial profile

$$u = 1 + \alpha_0 h_0(\sigma) + \alpha_2 h_2(\sigma),$$

where α_0 and $\alpha_2 > 0$ are constants to be chosen later, and $\sigma(s, t) = s/\sqrt{T-t}$. That is, we prescribe $\psi(s, t_0) = R_n(t_0) \cdot u(\sigma, t_0)$ as

$$(37) \quad \psi(s, t_0) = \sqrt{2(n-1)}\sqrt{T-t_0} \left[(1 + \alpha_0 - 2\alpha_2) + \alpha_2 \frac{s^2}{T-t_0} \right]$$

on the region $\mathcal{N} = \{s : |s| \leq s_0\}$, where s_0 is given by

$$s_0 = \varepsilon \sqrt{T-t_0} \log \frac{1}{T-t_0}.$$

We will require

$$(38) \quad -\varepsilon \leq \alpha_0 \leq 2\alpha_2$$

and

$$(39) \quad \frac{\varepsilon}{\tau_0} \leq \alpha_2 \leq \frac{1}{\tau_0},$$

where

$$\tau_0 = \log \frac{1}{T-t_0} \gg 1.$$

The radius of a neck of the form (37) is

$$(40) \quad r_0 = \psi(0, t_0) = \sqrt{2(n-1)}\sqrt{T-t_0}(1 + \alpha_0 - 2\alpha_2).$$

Thus

$$\psi = r_0 + \sqrt{2(n-1)} \frac{\alpha_2 s^2}{\sqrt{T-t_0}}.$$

Since

$$\psi_s = 2\sqrt{2(n-1)} \frac{\alpha_2 s}{\sqrt{T-t_0}},$$

we have $|\psi_s| \leq \psi_s(s_0, t_0)$ on $\mathcal{N}$, where $\psi_s(s_0, t_0) = 2\varepsilon\sqrt{2(n-1)} \cdot \tau_0\alpha_2$ satisfies

$$\varepsilon^2 \leq \frac{\psi_s(s_0, t_0)}{2\sqrt{2(n-1)}} \leq \varepsilon.$$

Outside $\mathcal{N}$, we extend ψ linearly with slope $\psi_s(s_0, t_0) > 0$ until we reach the height $\psi = 1$, where we glue on a standard profile that is smooth at the poles. As in §8 of [3], this can be done with $|\psi_s| \leq 1$ everywhere. Notice that the initial profile $\psi(\cdot, t_0)$ is smooth except on $\partial\mathcal{N}$; the corresponding solution of the Ricci flow will be smooth for all $t > t_0$ that it exists.

We next obtain a formula for the upper bound F^* on the cylindricity F of a neck of the form (37), where α_0 and α_2 satisfy (38) and (39), respectively. Note that $F \leq 0$ off $\mathcal{N}$. On $\mathcal{N}$, we have

$$\begin{aligned} \psi &\leq \psi(s_0, t_0) = r_0 + \sqrt{2(n-1)} \frac{\alpha_2 s_0^2}{\sqrt{T-t_0}} \\ &\leq \sqrt{2(n-1)}(1 + \varepsilon^2\tau_0)\sqrt{T-t_0} \end{aligned}$$

and

$$\psi_{ss} = 2\sqrt{2(n-1)} \frac{\alpha_2}{\sqrt{T-t_0}} \leq \frac{2\sqrt{2(n-1)}}{\tau_0\sqrt{T-t_0}}.$$

Independent of $\varepsilon \leq \frac{1}{4}$, we may choose t_* close enough to T so that for any $t_0 \in [t_*, T)$, the radius is bounded from below by

$$r_0 \geq \sqrt{2(n-1)}\sqrt{T-t_0} \left(1 - \varepsilon - \frac{2}{\tau_0}\right) \geq \frac{\sqrt{2(n-1)}}{2}\sqrt{T-t_0},$$

and so that the sectional curvature L is bounded from below on $\mathcal{N}$ by

$$L = \frac{1 - \psi_s^2}{\psi^2} \geq \frac{1 - 8(n-1)\varepsilon^2}{\psi(s_0, t_0)^2} \geq \frac{1}{4(n-1)(T-t_0)(1 + \varepsilon^2\tau_0)^2} \geq e^2,$$

provided $\tau_0 > \tau_0^*(n, \varepsilon)$. Here we used the fact that $\lim_{x \searrow 0} (x \log \frac{1}{x}) = 0$. By Lemma 9.1 of [3], the bound $L \geq e^2$ lets us assume $\lambda = 0$. Combining the estimates above, we conclude that there exists $C = C(n, t_*)$ such that for any $t_0 \in [t_*, T)$, one has

$$\begin{aligned} F &= \frac{K}{L} \log L = \frac{\psi\psi_{ss}}{1 - \psi_s^2} \log \frac{1 - \psi_s^2}{\psi^2} \\ &\leq \frac{2}{1 - 8(n-1)\varepsilon^2} \psi\psi_{ss} \log \frac{1}{r_0} \\ &\leq 16(n-1) \frac{1 + \varepsilon^2\tau_0}{\tau_0} (\log 2 + \frac{\tau_0}{2}) \\ &\leq C\varepsilon^2\tau_0. \end{aligned}$$

Hence we may define

$$(41) \quad F^* = (n-1) + C\varepsilon^2\tau_0.$$

This formula for F^* lets us adjust our constants in order to force pinching at T .

Lemma 21. *There exist ε_* and t_* depending only on n such that for any $\varepsilon \in (0, \varepsilon_*]$, all initial times $t_0 \in [t_*, T)$, and any α_2 satisfying (39), there exists some α_0 satisfying (38) such that the solution with initial profile (37) becomes singular exactly at T .*

Proof. Proposition 5.2 of [3] implies that r is monotonically decreasing. So for all $t \geq t_0$ that the solution exists, one has

$$r(t)^2 \leq r_0^2 \leq 2(n-1)(T-t_0).$$

As above, we fix $t_* < T$ so that (41) holds for all $t_0 \in [t_*, T)$. Then by (36), there exists $c > 0$ depending only on n such that

$$-2(n-1) \leq \frac{d}{dt}(r^2) \leq -2(n-1) + 2\frac{n-1+C\varepsilon^2\tau_0}{\tau_0-c}.$$

There exists $C_* = C_*(n, t_*)$ such that $\frac{n-1+C\varepsilon^2\tau_0}{\tau_0-c} \leq C_*\varepsilon^2$. Hence we may fix $\varepsilon_* = \varepsilon_*(C_*)$ such that for all $\varepsilon \in (0, \varepsilon_*]$, one has

$$-2(n-1) \leq \frac{d}{dt}(r^2) \leq -2(n-1)(1-\varepsilon).$$

By Lemma 7.1 of [3], the solution becomes singular at $\tilde{T}$ if and only if $r(\tilde{T}) = 0$. It follows that

$$\frac{r_0^2}{2(n-1)} \leq \tilde{T} - t_0 \leq \frac{1}{1-\varepsilon} \cdot \frac{r_0^2}{2(n-1)}.$$

By (40), this implies that

$$(1 + \alpha_0 - 2\alpha_2)^2 \leq \frac{\tilde{T} - t_0}{T - t_0} \leq \frac{(1 + \alpha_0 - 2\alpha_2)^2}{1 - \varepsilon}.$$

Thus if $\alpha_0 - 2\alpha_2 > 0$, then pinching occurs at $\tilde{T} > T$. On the other hand, if $-1 < \alpha_0 - 2\alpha_2 < -\varepsilon$, then pinching occurs at $\tilde{T} < T$. By Lemma 20, pinching will occur exactly at T for some $\alpha_0 \in [2\alpha_2 - \varepsilon, 2\alpha_2]$. Clearly, this α_0 satisfies (38). $\square$

2.17. The dominant asymptotic profile. For $\tau \geq \tau_0$, let $v(\sigma, \tau)$ denote the solution of the Ricci flow corresponding to the initial data (37). Then

$$v(\sigma, \tau_0) = \alpha_0 h_0(\sigma) + \alpha_2 h_2(\sigma) \quad \text{for} \quad |\sigma| \leq \sigma_0 = \sigma(s_0, t_0) = \varepsilon \tau_0.$$

For $\tau \geq \tau_0$, define $V(\sigma, \tau) = \eta(e^{-\delta\tau/2}\sigma)v(\sigma, \tau)$, noting that $V(\cdot, \tau) \in \mathcal{H}$ admits the orthogonal decomposition

$$(42) \quad V(\sigma, \tau) = a_0(\tau)h_0(\sigma) + a_2(\tau)h_2(\sigma) + W(\sigma, \tau).$$

Intuitively, one expects that a_0 and a_2 will be close to α_0 and α_2 , respectively, at $\tau = \tau_0$ and for a short time thereafter. We now make this expectation precise. Define $U \in \mathcal{H}$ by

$$U = V(\cdot, \tau_0) - \alpha_0 h_0 - \alpha_2 h_2.$$

At $\tau = \tau_0$, one has $|V_\sigma| = |v_\sigma| = |\psi_s|/\sqrt{2(n-1)} \leq \varepsilon$ for $|\sigma| \leq \sigma_0$ and $|V_\sigma| \leq 1$ everywhere. Because $|\alpha_0|, \alpha_2 \in [0, 1]$, there are universal constants C (which may change from line to line) such that

$$\begin{aligned} U(\sigma) &= 0 & \text{for all } |\sigma| < \sigma_0 \\ |U(\sigma)| &\leq C(1 + \sigma^2) & \text{for all } |\sigma| \geq \sigma_0, \end{aligned}$$

hence such that

$$\begin{aligned}\|U\|^2 &\leq C \int_{|\sigma| \geq \sigma_0} (1 + \sigma^2)^2 e^{-\sigma^2/4} d\sigma \\ &\leq C e^{-\sigma_0^2/5} = C e^{-\varepsilon^2 \tau_0^2/5}.\end{aligned}$$

Because $\|W\|^2 = \langle W, W \rangle = \langle W, U \rangle \leq \|W\| \|U\|$ at $\tau = \tau_0$, it follows that the constant W_0 introduced in Lemma 8 satisfies

$$(43) \quad W_0 = e^{\tau_0} \|W(\cdot, \tau_0)\| \leq e^{\tau_0} \|U\| \leq C \exp(\tau_0 - \frac{\varepsilon^2}{10} \tau_0^2).$$

Similarly, one has

$$a_2(\tau_0) = \frac{1}{\|h_2\|^2} \langle h_2, V(\cdot, \tau_0) \rangle = \alpha_2 + \frac{1}{\|h_2\|^2} \langle h_2, U \rangle,$$

and hence

$$(44) \quad |a_2(\tau_0) - \alpha_2| \leq C e^{-\varepsilon^2 \tau_0^2/10}.$$

We are now ready to prove that alternative (34) occurs. By (32), one has

$$\frac{d}{d\tau} a_2(\tau) = -8a_2(\tau)^2 + M^*(\tau),$$

where $|M^*(\tau)| \leq C(1+W_0)\tau^{-3}$. By (43), we may make τ_0 so large that $C(1+W_0) \leq C_0$. Then

$$-8a_2(\tau)^2 - \frac{C_0}{\tau^3} \leq \frac{d}{d\tau} a_2(\tau) \leq -8a_2(\tau)^2 + \frac{C_0}{\tau^3}.$$

By (39) and (44), we may make τ_0 larger if necessary so that

$$\sqrt{\frac{C_0}{8\tau_0^3}} \ll \frac{\varepsilon}{2\tau_0} \leq a_2(\tau_0) \leq \frac{2}{\tau_0}.$$

As long as $a_2(\tau)^2 > \frac{C_0}{8}\tau^{-3}$, the function $a_2(\tau)$ will be decreasing. For such times, one has

$$\frac{d}{d\tau} a_2(\tau) \geq -8a_2(\tau_0)^2 - \frac{C_0}{\tau^3} \geq -\frac{32}{\tau_0^2} - \frac{C_0}{\tau_0^3} \geq -\frac{64}{\tau_0^2}$$

provided that τ_0 is large enough, and hence

$$a_2(\tau) \geq a_2(\tau_0) - \frac{64}{\tau_0^2}(\tau - \tau_0) \geq \frac{\varepsilon}{2\tau_0} - \frac{64}{\tau_0^2}(\tau - \tau_0).$$

In particular, given any $C > 0$, one can make τ_0 so large that

$$a_2(\tau_0 + 1) \geq \frac{\varepsilon}{2\tau_0} - \frac{64}{\tau_0^2} \geq C \frac{1 + W_0}{(\tau_0 + 1)^2}.$$

This justifies our assumption that a_2 was decreasing for $\tau_0 \leq \tau \leq \tau_0 + 1$. Moreover, if we choose C as in (33) then it violates alternative (33). Hence we have proved the following result.

Proposition 22. *For any final time T , all sufficiently small $\varepsilon > 0$, all initial times t_0 sufficiently close to T , and any α_2 satisfying (39), there exists α_0 satisfying (38) such that the solution (42) with initial profile (37) becomes singular at T and satisfies the asymptotic profile*

$$(45) \quad a_2(\tau) = \frac{1}{8\tau} + o\left(\frac{1}{\tau}\right) \quad \text{as} \quad \tau \rightarrow \infty.$$

2.18. The intermediate region. Consider a solution that forms a singularity at time $t = T$. Define u, v , etc. as above and assume that our solution satisfies (45).

We have shown that

$$(46) \quad v(\sigma, \tau) = \frac{\sigma^2 - 2}{8\tau} + o\left(\frac{1}{\tau}\right) \quad (\tau \rightarrow \infty),$$

uniformly on any bounded interval $|\sigma| \leq A$.

Now consider the quantity

$$q = u^2 - 1.$$

Lemma 23. *For any constants $\varepsilon > 0$ and $B < \infty$, there exist A and $\bar{\tau}$ such that*

$$(1 - \varepsilon) \frac{\sigma^2}{4\tau} < q(\sigma, \tau) < (1 + \varepsilon) \frac{\sigma^2}{4\tau}$$

whenever $A \leq \sigma \leq B\sqrt{\tau}$ and times $\tau \geq \bar{\tau}$.

In other words, for $1 \ll \sigma \leq B\sqrt{\tau}$ one has

$$u = \sqrt{1 + (1 + o(1)) \frac{\sigma^2}{4\tau}} \approx \sqrt{1 + \frac{\sigma^2}{4\tau}}.$$

We shall prove this Lemma by computing the evolution equation for q in the variables τ and $\rho = \sigma/\sqrt{\tau}$, and finding suitable sub- and supersolutions. The evolution equation (49) turns out to be a small perturbation of a first order linear equation, which is easily solved by following its characteristics.

2.18.1. Equation for q . Since $q_\sigma = 2uu_\sigma$ and $q_{\sigma\sigma} = 2uu_{\sigma\sigma} + 2u_\sigma^2$, one has

$$(47) \quad q_\tau|_x = q_{\sigma\sigma} + q + \frac{n-2}{2} \frac{q_\sigma^2}{q+1}.$$

Here the time derivative is with constant x . Using Lemma 5, we get

$$(48) \quad q_\tau|_\sigma = q_{\sigma\sigma} - \frac{\sigma}{2} q_\sigma + q - nJ(\sigma, \tau)q_\sigma + \frac{n-2}{2} \frac{q_\sigma^2}{q+1},$$

where $q_\tau|_\sigma$ is the derivative of q with σ held constant and where

$$J = \frac{u_\sigma}{u} + \int_0^\sigma \frac{u_\sigma^2}{u^2} d\sigma.$$

We now regard q as a function of the new variable $\rho = \sigma/\sqrt{\tau}$. Because

$$q_\tau|_\sigma = q_\tau|_\rho - \frac{1}{2} \frac{\rho}{\tau} q_\rho,$$

one has

$$(49) \quad q_\tau|_\rho - \frac{1}{\tau} q_{\rho\rho} + \frac{\rho}{2} q_\rho - q = \frac{1}{\tau} \left\{ \frac{\rho}{2} q_\rho - n\sqrt{\tau} J q_\rho + \frac{n-2}{2} \frac{q_\rho^2}{q+1} \right\}.$$

2.18.2. *A subsolution for (49).* We begin with some simple estimates. Fix some $A_0 \gg 1$ and recall that τ_0 is the smallest time under consideration in this paper. Given $B > 1$, we shall choose $A \geq A_0$ and $\tau_1 \geq \tau_0$ large enough to make everything work. The region $A/\sqrt{\tau} \leq \rho \leq B^2$ corresponds to $A \leq \sigma \leq B^2\sqrt{\tau}$. By Lemma 4, there exist $C_0 = C_0(B)$ and $C_1 = C_1(A_0)$ such that for all $A \leq \sigma \leq B^2\sqrt{\tau}$, one has

$$u \leq 1 + C_0 \frac{\sigma^2}{\tau} = 1 + C_0 \rho^2 \leq 1 + C_0 B^4$$

and

$$0 \leq u_\sigma \leq C_0 \frac{1 + \sigma}{\tau} \leq C_1 \frac{\sigma}{\tau} = C_1 \frac{\rho}{\sqrt{\tau}}.$$

Thus we have

$$0 \leq u_\rho \leq C_1 \rho$$

and

$$0 \leq \rho q_\rho = 2\rho u_\rho u \leq C(A_0, B)$$

for $A/\sqrt{\tau} \leq \rho \leq B^2$. Since $u \geq 1 - C/\tau$ for $C = C(n, F^*)$, we may assume without loss of generality that $\tau_0 \gg 1$ is so large that

$$\frac{q_\rho^2}{q+1} = \frac{q_\rho^2}{u^2} \leq 2q_\rho^2 \leq C(A_0, B)$$

for $A/\sqrt{\tau} \leq \rho \leq B^2$ and $\tau \geq \tau_0$. Since $n\sqrt{\tau}Jq_\rho \geq 0$ for all $0 \leq \sigma \leq e^{\delta\tau}$, it follows that

$$(50) \quad \frac{\rho}{2}q_\rho - n\sqrt{\tau}Jq_\rho + \frac{n-2}{2} \frac{q_\rho^2}{q+1} \leq C(A_0, B).$$

We introduce the linear differential operator

$$\mathcal{L}[f] \doteq f_\tau|_\rho - \frac{1}{\tau}f_{\rho\rho} + \frac{\rho}{2}f_\rho - f.$$

Then (49) and (50) imply that $q = u^2 - 1$ satisfies

$$|\mathcal{L}[q]| \leq \frac{C(A_0, B)}{\tau}.$$

whenever $A/\sqrt{\tau} \leq \rho \leq B^2$ and $\tau \geq \tau_0$.

Let Q be a smooth function with support in $[0, 1]$ such that $Q(r) \leq r^2/4$, and define

$$K_0 = \max_{0 \leq r \leq 1} |Q''(r)|.$$

For $\tau_1 \geq \tau_0$, define

$$(51) \quad q_0(\rho, \tau) = \rho_1(\tau)^2 Q(\rho/\rho_1(\tau)),$$

with

$$\rho_1(\tau) \doteq e^{(\tau-\tau_1)/2} \frac{A}{\sqrt{\tau_1}}.$$

This function is almost in the nullspace of the linear operator $\mathcal{L}$; indeed, a straightforward computation shows that

$$\mathcal{L}[q_0] = -\frac{1}{\tau} \frac{\partial^2 q_0}{\partial \rho^2} = -\frac{1}{\tau} Q''(\rho/\rho_1(\tau)),$$

so that $|\mathcal{L}[q_0]| \leq K_0/\tau$.

Now for $K \geq K_0$ to be determined and any $\theta \in (1/2, 1)$, consider

$$(52) \quad q_-(\rho, \tau) = \theta q_0(\rho, \tau) + \frac{K}{\tau}.$$

One then has

$$\mathcal{L}[q_-] = \theta \mathcal{L}[q_0] + K \mathcal{L}[\tau^{-1}] < \theta \frac{K_0}{\tau} - \frac{K}{\tau} < \frac{K_0 - K}{\tau} \leq 0,$$

so that q_- is a subsolution for all $\tau \geq \tau_0$. Moreover, if we choose $K \geq K_0 + C(A_0, B)$, then

$$(53) \quad \mathcal{L}[q_- - q] \leq \frac{C(A_0, B) + K_0 - K}{\tau} \leq 0$$

for $A/\sqrt{\tau} \leq \rho \leq B^2$ and $\tau \geq \tau_0$.

Define the region

$$(54) \quad \Omega = \left\{ (\rho, \tau) : \frac{A}{\sqrt{\tau}} \leq \rho \leq \rho_1(\tau), \tau_1 \leq \tau \leq \omega \right\},$$

where ω is determined by $\rho_1(\omega) = B^2$, i.e.

$$\omega = \tau_1 + \log(B^4 \tau_1 / A^2).$$

We will choose A and τ_1 so that $B^4 \tau_1 > A^2$. First, we compare the values of q and q_- along the parabolic boundary of Ω .

Along the left edge $\rho = A/\sqrt{\tau}$, $\tau \geq \tau_1$, it follows from (46) that

$$(55) \quad q(A/\sqrt{\tau}, \tau) = u^2(A, \tau) - 1 = \frac{A^2 - 2}{4\tau} + o\left(\frac{1}{\tau}\right) \geq (1 - \varepsilon_0) \frac{A^2}{4\tau},$$

where $\varepsilon_0 = \varepsilon_0(A_0, \tau_1)$. On the other hand, there exists $\varepsilon_1 = \varepsilon_1(A, K, \tau_0)$ such that

$$q_-(\rho, \tau) \leq \frac{\theta}{4} \rho^2 + \frac{K}{\tau} \leq (1 + \varepsilon_1) \frac{\theta}{4} \rho^2$$

for $\rho \geq A/\sqrt{\tau}$. So $q_- < q$ at $(A/\sqrt{\tau}, \tau)$ provided that

$$\frac{1}{2} \leq \theta < \frac{1 - \varepsilon_0}{1 + \varepsilon_1}.$$

Along the right edge $\rho = \rho_1(\tau)$, one has

$$q_-(\rho_1(\tau), \tau) = \rho_1(\tau)^2 Q(1) + \frac{K}{\tau} = \frac{K}{\tau},$$

because Q is supported in $[0, 1]$. But since $q_\rho \geq 0$, estimate (55) implies that

$$q(\rho_1(\tau), \tau) = q(e^{(\tau - \tau_1)/2} A / \sqrt{\tau_1}, \tau) \geq q(A/\sqrt{\tau}, \tau) \geq \frac{1 - \varepsilon_0}{4} \frac{A^2}{\tau},$$

where $\varepsilon_0 = \varepsilon_0(A_0, \tau_1)$. So $q_- \leq q$ at $(\rho_1(\tau), \tau)$ provided that

$$A \geq 2\sqrt{\frac{K}{1 - \varepsilon_0}}.$$

The bottom of Ω is the single point $(A/\sqrt{\tau_1}, \tau_1)$. So the choice $\theta < (1 - \varepsilon_0)/(1 + \varepsilon_1)$ ensures that $q_- < q$ there as well.

Now choose K depending on A_0 , B , and K_0 to satisfy $K \geq K_0 + C(A_0, B)$. Take $\tau_1 \geq \tau_0$ large enough so that $\varepsilon_0 < 1/2$ and choose $A \geq A_0$ depending on K so that $A \geq \sqrt{8K}$ and so that $\varepsilon_1 > 0$ is as small as desired. Then increase τ_1 if necessary

so that $\tau_1 \geq A^2/B^4$, making ε_0 as small as desired. These choices make it possible to take $\theta \in (1/2, 1)$ as close to 1 as one wishes and still satisfy $\theta < (1 - \varepsilon_0)/(1 + \varepsilon_1)$. Furthermore, they ensure that $q_- \leq q$ along the parabolic boundary of Ω . Because $e^{(\tau - \tau_1)/2} A/\sqrt{\tau_1} \leq B^2$ for all $\tau \leq \omega$, it follows from (53) that $q \geq q_-$ throughout Ω . Since for any ε , one may take $Q(r) = r^2/4$ for $r \in (\varepsilon, 1 - \varepsilon)$, this implies in particular that

$$q(\rho, \tau) > \theta \frac{\rho^2}{4}$$

for all points ρ such that $A/\sqrt{\tau} \leq \rho \leq B$ and times τ such that

$$2 \log(B\sqrt{\tau_1}/A) \leq \tau - \tau_1 \leq 2[\log(B\sqrt{\tau_1}/A) + \log B].$$

Our estimates only improve if we increase τ_1 (without altering any other choices); in particular, θ does not decrease. Since $\tau = \tau_1 + o(\tau_1)$ as $\tau_1 \rightarrow \infty$, we have proved the lower estimate in Lemma 23.

2.18.3. A supersolution. A slight modification of the preceding construction will give an upper barrier for q on the same domain Ω . One begins with the *a priori* bound

$$(56) \quad u \leq C(1 + \sigma^2/\tau) = C(1 + \rho^2) \text{ and hence } q \leq C(1 + \rho^2)\rho^2 = C_0(B)\rho^2,$$

which holds for $A_0/\sqrt{\tau} \leq \rho \leq B^2$.

Choose a smooth function $\hat{Q}$ which satisfies $\hat{Q}(r) \geq r^2/4$, and define $\hat{K}_0 = \max_{[0,1]} |\hat{Q}''|$. Instead of requiring $\hat{Q}$ to be supported in $[0, 1]$, we impose

$$\hat{Q}(1) = 2C_0(B) \text{ with } C_0(B) \text{ as in (56).}$$

For $\tau \geq \tau_1$, one introduces $\hat{q}_0(\rho, \tau) = \rho_1(\tau)^2 \hat{Q}(\rho/\rho_1(\tau))$, as in (51), and finds that $|\mathcal{L}[\hat{q}_0]| \leq \hat{K}_0/\tau$. For $\hat{K} \geq \hat{K}_0$ and any $\theta \in (1, \frac{3}{2})$, define

$$q_+(\rho, \tau) = \theta \hat{q}_0(\rho, \tau) - \hat{K}/\tau.$$

Choosing $\hat{K} \geq \hat{K}_0 + C(A_0, B)$, one finds as in (53) that $\mathcal{L}[q_+ - q] \geq 0$ for $\tau \geq \tau_0$ and $A/\sqrt{\tau} \leq \rho \leq B^2$. Thus q_+ is a supersolution, provided $q_+ \geq q$ on the parabolic boundary of Ω .

On the left edge of Ω , one has, as in (55), $q \leq (1 + \varepsilon_0)A^2/4\tau$. One also has $q_+ \geq (1 - \varepsilon_1)\theta A^2/4\tau$, where $\varepsilon_1 = \varepsilon_1(A, K)$.

Hence one will have $q \leq q_+$ on the left edge if θ satisfies

$$\frac{1 + \varepsilon_0}{1 - \varepsilon_1} < \theta < \frac{3}{2}.$$

On the right edge of Ω one has $\rho = \rho_1(\tau)$, and thus

$$q_+(\rho, \tau) = \theta \rho_1(\tau)^2 \hat{Q}(1) - \hat{K}/\tau \geq \theta(1 - \hat{K}/A^2)Q(1)\rho_1(\tau)^2 Q(1)\rho_1(\tau)^2,$$

provided A is chosen so large that $\theta(1 - \hat{K}/A^2) > 1$. On the other hand we know that $q \leq C_0(B)\rho_1(\tau)^2$ holds on the right edge of Ω . Hence our condition $\hat{Q}(1) > C_0(B)$ implies that $q_+ > q$ on the right edge.

It now follows that $q \leq q_+$ on Ω , and as with the subsolution, this implies that $q(\rho, \tau) < \theta \rho^2/4$ for all $\rho \in [A/\sqrt{\tau}, B]$ and $\tau \geq \tau_1 + o(\tau_1)$.

2.19. Convergence in the outer region. Let B and $\varepsilon > 0$ be given constants. Define $A = A(\varepsilon, B)$ and $\bar{\tau} = \bar{\tau}(\varepsilon, B)$ as in Lemma 23. Then we have shown that when

$$(57) \quad \sigma = B\sqrt{\tau}, \text{ i.e. when } s^2 = \frac{B^2}{4}(T-t) \log \frac{1}{T-t},$$

one has

$$\sqrt{1 + (1 - \varepsilon)B^2/4} \leq u \leq \sqrt{1 + (1 + \varepsilon)B^2/4}.$$

For $\psi = R_n(t)u$, this therefore means that

$$(58) \quad \sqrt{T-t} \sqrt{1 + (1 - \varepsilon)B^2/4} \leq \frac{\psi}{\sqrt{2(n-1)}} \leq \sqrt{T-t} \sqrt{1 + (1 + \varepsilon)B^2/4}$$

for all $t \in [\bar{t}, T)$, where $\bar{t} = T - e^{-\bar{\tau}}$.

We define the *outer region* (with parameter B) to be the portion of space-time given by

$$\Gamma_B = \{(x, t) : \bar{\tau} \leq \tau < T, s(x, t)^2 \geq \frac{B^2}{4}(T-t) \log \frac{1}{T-t}\}.$$

The boundary $\partial\Gamma_B$ of Γ_B is given by (57).

For any given $t_* \in [\bar{t}, T)$, we let x_* be the x coordinate of the point on $\partial\Gamma_B$ with $t = t_*$. We will now show that $s(x_*, t)$ and $\psi(x_*, t)$ change very little for $t_* \leq t < T$. This will directly imply the asymptotic description of $\psi(s, T)$ for $s \rightarrow 0$ in the introduction. Fix some $B_0 > 1$. We begin with a simple uniform estimate for ψ_s in the region $B_0\sqrt{\tau} \leq \sigma \leq \eta e^{\delta\tau}$.

Lemma 24. *For any $B_0 > 1$, there exist constants $\gamma, \delta > 0$ and $C < \infty$ depending only on B_0 and the initial data such that*

$$0 \leq \psi_s(s, t) \leq \frac{C}{\sqrt{\tau}}$$

for all $\tau \geq \bar{\tau}(B_0)$ and $B_0\sqrt{\tau} \leq \sigma(s) \leq \gamma e^{\delta\tau}$.

Proof. Define $r(t) = \psi(0, t)$. By (45), we have

$$r = \sqrt{2(n-1)(T-t)} \left\{ 1 - \frac{1}{4\tau} + o\left(\frac{1}{\tau}\right) \right\}.$$

By (58), we have

$$\psi(s, t) \geq \psi(\bar{s}, t) \geq \sqrt{2(n-1)(T-t)} \sqrt{1 - B_0^2/8}$$

for all $s \geq \bar{s}(B_0) = B_0\tau\sqrt{T-t}$ and $\tau \geq \bar{\tau}(B_0) \geq \tau_0$. Now restricting our attention to the region of the neck where $\psi \leq 1/B_0$ and $0 \leq \psi_s \leq 1/B_0$, we may apply estimate (34) of [3], obtaining C depending on B_0 and F^* such that

$$\psi_s^2 \leq C \log \left(\frac{\log r}{\log \psi} \right).$$

This estimate holds as long as $\sigma \leq \gamma e^{\delta\tau}$, i.e. as long as $s \leq \gamma(\sqrt{T-t})^{1-2\delta} \leq r^{1-2\delta}$. Let $c_1 = \log \sqrt{1 - B_0^2/8}$. Then

$$\frac{\log r}{\log \psi} \leq \frac{\log \sqrt{2(n-1)} - \frac{\tau}{2} - \frac{1}{4\tau} + o(\frac{1}{\tau})}{\log \sqrt{2(n-1)} - \frac{\tau}{2} + c_1} = 1 + \frac{2c_1}{\tau} + o(\frac{1}{\tau}).$$

Hence $\tau\psi_s^2 \leq C\tau \log\left(\frac{\log r}{\log \psi}\right) = C(2c_1 + o(1))$. $\square$

We are now ready to describe $s(x_*, t)$ and $\psi(x_*, t)$.

Lemma 25. *For any x_* the limits*

$$\lim_{t \rightarrow T} s(x_*, t) \text{ and } \lim_{t \rightarrow T} \psi(x_*, t)$$

exist. Furthermore, a constant $C < \infty$ which does not depend on $B \geq B_0$ or x_ exists for which one has*

$$(59) \quad |s(x_*, t) - s(x_*, t_*)| \leq C\{B^{-3} + |\log(T - t_*)|^{-3/2}\} s(x_*, t_*)$$

and


$$(60) \quad |\psi(x_*, t) - \psi(x_*, t_*)| \leq \frac{C}{B^2} \psi(x_*, t_*)$$

for all $t \in [t_*, T)$.

Proof. To estimate the change in $s(x_*, t)$ we use

$$(61) \quad \left. \frac{\partial s}{\partial t} \right|_{x=x_*} = n \frac{\psi_s(x_*, t)}{\psi(x_*, t)} + n \int_0^{s(x_*, t)} \left(\frac{\psi_s}{\psi} \right)^2 ds,$$

as in the proof of Lemma 2. Let $t' \in (t_*, T)$ be given, and let x' be such that $s(x', t')$ lies on $\partial\Gamma_B$.



Then at time t' , Lemma 24 implies that

$$(62) \quad 0 \leq \psi_s(s, t') \leq \frac{C}{\sqrt{\log \frac{1}{T-t'}}} \text{ for all } s \leq s'.$$

Since $x_* \geq x'$, we then get

$$\int_0^{s(x_*, t')} \left(\frac{\psi_s}{\psi} \right)^2 ds = \int_0^{s(x', t')} \left(\frac{\psi_s}{\psi} \right)^2 ds + \int_{s(x', t')}^{s(x_*, t')} \left(\frac{\psi_s}{\psi} \right)^2 ds.$$

In the first term, we use (62) to get

$$\begin{aligned} \int_0^{s(x', t')} \left(\frac{\psi_s}{\psi} \right)^2 ds &\leq \frac{C}{\sqrt{\ln 1/(T - t_*)}} \int_0^{s(x', t')} \frac{\psi_s}{\psi^2} ds \\ &\leq \frac{C}{\sqrt{\ln 1/(T - t_*)}} \frac{1}{\psi(0, t')} \\ &\leq \frac{C}{\sqrt{(T - t') \ln 1/(T - t_*)}}. \end{aligned}$$

In the second term, we have $x' < x < x_*$ so that

$$\psi \geq \psi(x', t') \geq \sqrt{2(n-1)(T-t')(1+(1-\varepsilon)B^2/4)} \geq CB\sqrt{T-t'},$$

with C independent of t or $B \geq B_0$. On the other hand, we also have $\psi_s \leq 1$, so that

$$\begin{aligned} \int_{s(x', t')}^{s(x_*, t')} \left(\frac{\psi_s}{\psi} \right)^2 ds &\leq \int_{s(x', t')}^{s(x_*, t')} \left(\frac{\psi_s}{\psi^2} \right) ds \\ &\leq \frac{1}{\psi(x', t')} \\ &\leq \frac{C}{B\sqrt{T-t'}}. \end{aligned}$$

For the first term in (61), one has the same estimate, so that at any time $t' \in (t_*, T]$, one has

$$0 \leq \partial_t s(x_*, t') \leq C\{B^{-1} + |\log(T - t_*)|^{-1/2}\}(T - t')^{-1/2}.$$

This is integrable in $t' \in (t_*, T)$, so that $\lim_{t \rightarrow T} s(x_*, t)$ must exist. Integration with $t_* < t' < t$ then leads to

$$|s(x_*, t) - s(x_*, t_*)| \leq C\{B^{-1} + |\log(T - t_*)|^{-1/2}\}\sqrt{T - t_*}.$$

Recall (57), apply $a^2b, ab^2 \leq a^3 + b^3$ to $a = B^{-1}$ and $b = |\log(T - t_*)|^{-1/2}$, and conclude

$$\begin{aligned} |s(x_*, t) - s(x_*, t_*)| &\leq C\{B^{-1} + |\log(T - t_*)|^{-1/2}\} \frac{s(x_*, t_*)}{B\sqrt{\ln 1/(T - t_*)}} \\ &\leq C\{B^{-3} + |\log(T - t_*)|^{-3/2}\} s(x_*, t_*), \end{aligned}$$

which proves the estimate (59) for s .

To establish the other estimate, we recall from Proposition 5.3 in [3] that there is a constant C which only depends on the initial data and the dimension n such that

$$|2\psi\psi_t| \leq C$$

always holds. In particular, the constant C does not depend on our choice of B . In [3], we used this estimate to show that $\lim_{t \rightarrow T} \psi(x_*, t)$ exists. It also implies that

$$(63) \quad |\psi(x_*, t)^2 - \psi(x_*, t_*)^2| \leq C(t - t_*)$$

and hence, using (58), that

$$|\psi(x_*, t) - \psi(x_*, t_*)| \leq \frac{C(t - t_*)}{\psi(x_*, t) + \psi(x_*, t_*)} \leq \frac{C}{B} \sqrt{T - t_*} \leq \frac{C}{B^2} \psi(x_*, t_*)$$

for all $t \in (t_*, T)$. $\square$

2.20. Asymptotics in the outer region. Using (58) and (60), we compute that

$$\begin{aligned}\psi(x_*, t) &= (1 + \mathcal{O}(B^{-2}))\psi(x_*, t_*) \\ &= (1 + \mathcal{O}(B^{-2}))\sqrt{2(n-1)[1 + (1 + \theta\varepsilon)B^2/4]}(T - t_*) \\ &= (1 + \mathcal{O}(B^{-2} + \varepsilon))\sqrt{\frac{n-1}{2}}B\sqrt{T - t_*}\end{aligned}$$

for some $|\theta| \leq 1$.

From the definition (57), we see that $s = s(x_*, t_*)$ satisfies

$$B\sqrt{T - t} = \frac{s}{\sqrt{-\log(T - t)}}.$$

Hence, taking logarithms and abbreviating $\omega = T - t$,

$$\log B + \frac{1}{2} \ln \omega = \log s - \frac{1}{2} \ln |\log \omega|,$$

so that

$$\begin{aligned}\log \omega &= 2 \ln s - \log |\log \omega| - 2 \ln B = \mathcal{O}(\log s) \\ &= 2 \ln s + \mathcal{O}(\log |\log s|) \\ &= (2 + o(1)) \log s\end{aligned}$$

We therefore find that

$$B\sqrt{\omega} = (1 + o(1))\frac{s}{\sqrt{-2 \ln s}}.$$

Thus as $t_* \rightarrow T$, one has

$$\psi = (1 + \mathcal{O}(B^{-2} + \varepsilon))\left(\frac{1}{2}\sqrt{n-1} + o(1)\right)\frac{s}{\sqrt{-\log s}}$$

for all $t \in [t_*, T]$ and x_* for which $s = s(x_*, t_*)$ satisfies (57). Since B and ε can be chosen arbitrarily, this implies

Lemma 26. *For any $\delta > 0$ there exist $B > 0$ and $\bar{t} < T$ such that*

$$\left(\frac{1}{2}\sqrt{n-1} - \delta\right)\frac{s}{\sqrt{-\log s}} \leq \psi \leq \left(\frac{1}{2}\sqrt{n-1} + \delta\right)\frac{s}{\sqrt{-\log s}}$$

in the region $(T - t)|\log(T - t)| \leq s^2/B^2$, $\bar{t} \leq t \leq T$.

3. FORMAL ASYMPTOTICS FOR GENERAL NECKPINCHES

The usual starting point for obtaining matched asymptotic expansions near a stationary solution of a nonlinear PDE is to linearize around that solution. This method encounters two obstacles when applied to the Ricci flow.

The first challenge is the well-known fact that the Ricci flow is only weakly parabolic. This deficiency stems from its invariance under the full diffeomorphism group of the underlying manifold. We overcome it by a variant of the DeTurck trick [9, 10]. (See § 3.4.) Our choice is equivalent to the Bianchi gauge, as adopted in the elliptic context by Biquard [6] and others.

The second challenge is more fundamental: it is the fact that the linearized PDE possesses a null eigenvalue, suggesting the presence of a center manifold. This phenomenon also occurs in neck pinches of the mean curvature flow [5], and flat self-similar solutions to the reaction diffusion equation $\partial_t u = \Delta u + u^p$ [20, 21, 22, 13]. We overcome it by carrying out a higher-order variational analysis. A heuristic description of our method is as follows. We will formally compute the dynamics of a center manifold of a fixed point of a flow. That fixed point is the *cylinder soliton* introduced in § 3.1, and the flow is the *dilated Ricci flow* introduced in § 3.2. The center manifold will be tangential to the null eigenspace of the linearization obtained in § 3.4. A general point on the center manifold will be of the form $\zeta + \Phi(\zeta)$, where ζ belongs to the kernel of the linearization and $\Phi(\zeta)$ is at least quadratic in its dependence on ζ . (See § 3.9.) To compute the formal dynamics, we substitute this *Ansatz* into the dilated Ricci flow, project onto the kernel of the linearization, and compute the purely quadratic terms, yielding an ODE on the kernel. The errors introduced by this method should be smaller than the order of the solution, namely $\mathcal{O}(\tau^{-2})$, where τ is the time scale introduced above in Section 2.

We pursue this method below. The computations it requires are extensive, but the (formal) conclusion it yields is eminently simple. It tells us that the asymptotics obtained rigorously in Section 2 for rotationally-symmetric solutions should be stable for fully general solutions.

3.1. The cylinder soliton. Recall that a *Ricci soliton* is a tuple $(\mathcal{M}^m, g, X)$, where $\mathcal{M}^m$ is a smooth manifold, g is its Riemannian metric, and X is a complete vector field on $\mathcal{M}^m$ such that the identity

$$-2 \operatorname{Rc}(g) = \mathcal{L}_X g + \lambda g$$

holds for some $\lambda \in \{-1, 0, 1\}$. It is well known that each Ricci soliton gives rise to a *self-similar solution* $\bar{g}(t)$ of the Ricci flow

$$\frac{\partial}{\partial t} \bar{g} = -2 \operatorname{Rc}(\bar{g})$$

defined by

$$\bar{g}(t) = (T + \lambda t) (\varphi_t^* g),$$

where φ_t is the one-parameter family of diffeomorphisms generated by the vector fields $(T + \lambda t)^{-1} X$.

Now consider the manifold $\mathbb{R} \times S^n$ with local coordinates $(y^0; y^1, \dots, y^n)$, where $y^0 \equiv x \in \mathbb{R}$ and $(y^1, \dots, y^n) \equiv y \in S^n$. Let $\hat{g} \equiv g_{\text{can}}$ denote the round metric of unit radius on S^n . Define a product metric g on $\mathbb{R} \times S^n$ by

$$(64) \quad g = dx \otimes dx + 2(n-1) \hat{g},$$

noting that its Ricci curvature satisfies

$$\operatorname{Rc}(g) = (n-1) \hat{g}.$$

Let X be the vector field on $\mathbb{R} \times S^n$ defined by

$$(65) \quad X = \operatorname{grad} \left(\frac{x^2}{4} \right) = \frac{x}{2} \frac{\partial}{\partial x},$$

noting that the Lie derivative of g with respect to X is

$$\mathcal{L}_X g = dx \otimes dx.$$

It is then easy to see that

$$(66) \quad -2 \operatorname{Rc}(g) = \mathcal{L}_X g - g,$$

hence that $(\mathbb{R} \times S^n, g, X)$ is a shrinking gradient Ricci soliton. We call this the *cylinder soliton*.

For later use, we observe that the Levi-Civita connection of g is given in local coordinates $(y^0; y^1, \dots, y^n)$ by the Christoffel symbols

$$(67) \quad \Gamma_{ij}^k = \begin{cases} \hat{\Gamma}_{ij}^k & \text{if } 1 \leq i, j, k \leq n \\ 0 & \text{otherwise.} \end{cases}$$

Here and throughout the remainder of this paper, a hat designates a geometric quantity associated to the round unit sphere $(S^n, \hat{g})$. The components $R_{ijk\ell} = g_{\ell m} R_{ijk}^m$ of the Riemannian curvature of g are

$$(68) \quad R_{ijk\ell} = \begin{cases} 2(n-1) \hat{R}_{ijk\ell} = 2(n-1) (\hat{g}_{i\ell} \hat{g}_{jk} - \hat{g}_{ik} \hat{g}_{j\ell}) & \text{if } 1 \leq i, j, k, \ell \leq n \\ 0 & \text{otherwise,} \end{cases}$$

and those of its Ricci tensor are

$$(69) \quad R_{ij} = \begin{cases} \hat{R}_{ij} = (n-1) \hat{g}_{jk} & \text{if } 1 \leq i, j \leq n \\ 0 & \text{otherwise.} \end{cases}$$

3.2. The dilated Ricci flow. Motivated by the rigorous results obtained in Section 2, we want to study finite-time local singularities of the Ricci flow that are modeled on the cylinder soliton. Accordingly, let $(\mathcal{M}^m, G(t))$ be a solution of the Ricci flow

$$\frac{\partial}{\partial t} G = -2 \operatorname{Rc}(G)$$

that exists for $0 \leq t < T < \infty$. Given any vector field X on $\mathcal{M}^m$, let φ_t be the family of diffeomorphisms solving

$$\frac{\partial}{\partial t} \varphi(\cdot, t) = \frac{1}{T-t} X(\varphi(\cdot, t)).$$

Then the *blow-up of G at T modified by X* is the metric $g(t)$ defined by

$$(70) \quad G(t) = (T-t) (\varphi_t^* g(t)).$$

Observing that

$$\frac{\partial}{\partial t} (\varphi_t^* g) = \frac{1}{T-t} \varphi_t^* (\mathcal{L}_X g) + \varphi_t^* \left(\frac{\partial}{\partial t} g \right),$$

one computes that

$$\varphi_t^* (-2 \operatorname{Rc}(g)) = -2 \operatorname{Rc}(G) = \frac{\partial}{\partial t} G = \varphi_t^* \left(-g + \mathcal{L}_X g + (T-t) \frac{\partial}{\partial t} g \right).$$

Thus if one regards g as a function of the rescaled time variable

$$(71) \quad \tau = \log \frac{1}{T-t},$$

then $(\mathcal{M}^m, g(\tau), X)$ becomes a solution of the *dilated Ricci flow*:

$$(72) \quad \frac{\partial}{\partial \tau} g = -2 \operatorname{Rc}(g) - \mathcal{L}_X g + g.$$

The following is an immediate consequence of equations (66) and (72).

Lemma 27. *The cylinder soliton $(\mathbb{R} \times S^n, g, X)$ is a stationary solution of the dilated Ricci flow.*

3.3. First and second order variation formulas. If h is a symmetric $(2, 0)$ -tensor on a Riemannian manifold $(\mathcal{M}^m, g)$, its Lichnerowicz Laplacian $\Delta_\ell h$ is defined by

$$(73) \quad \Delta_\ell h_{ij} = \Delta h_{ij} + 2R_{ipqj}h^{pq} - R_i^k h_{kj} - R_j^k h_{ik},$$

where Δh is the rough Laplacian. Its divergence δh is defined by

$$(74) \quad (\delta h)_j = -\nabla^i h_{ij}.$$

The following variation formulas are proved by direct calculation.

Lemma 28. *Let $(\mathcal{M}^m, g)$ be a Riemannian manifold; let h be a symmetric $(2, 0)$ -tensor on $\mathcal{M}^m$; and let ε be small enough that*

$$\tilde{g} = g + \varepsilon h$$

is a Riemannian metric.

- (1) *In local coordinates, the Christoffel symbols $\tilde{\Gamma}$ of the Levi-Civita connection of $\tilde{g}$ are*

$$\begin{aligned} \tilde{\Gamma}_{ij}^k &= \Gamma_{ij}^k + \frac{\varepsilon}{2} (\nabla_i h_j^k + \nabla_j h_i^k - \nabla^k h_{ij}) \\ &\quad - \frac{\varepsilon^2}{2} h_\ell^k (\nabla_i h_j^\ell + \nabla_j h_i^\ell - \nabla^\ell h_{ij}) + \mathcal{O}(\varepsilon^3). \end{aligned}$$

- (2) *In local coordinates, the Riemann tensor $\widetilde{\text{Rm}}$ of $\tilde{g}$ is*

$$\begin{aligned} \tilde{R}_{ijk}^\ell &= R_{ijk}^\ell + \frac{\varepsilon}{2} \left(\nabla_i \nabla_k h_j^\ell - \nabla_i \nabla^\ell h_{jk} - \nabla_j \nabla_k h_i^\ell + \nabla_j \nabla^\ell h_{ik} \right. \\ &\quad \left. + R_{ijm}^\ell h_k^m - R_{ijk}^m h_m^\ell \right) \\ &\quad + \frac{\varepsilon^2}{2} h_m^\ell (\nabla_i \nabla^m h_{jk} - \nabla_j \nabla^m h_{ik} - \nabla_i \nabla_k h_j^m + \nabla_j \nabla_k h_i^m) \\ &\quad + \frac{\varepsilon^2}{4} (\nabla_i h_k^m + \nabla_k h_i^m - \nabla^m h_{ik}) (\nabla_j h_m^\ell + \nabla^\ell h_{jm} - \nabla_m h_j^\ell) \\ &\quad - \frac{\varepsilon^2}{4} (\nabla_j h_k^m + \nabla_k h_j^m - \nabla^m h_{jk}) (\nabla_i h_m^\ell + \nabla^\ell h_{im} - \nabla_m h_i^\ell) \\ &\quad + \frac{\varepsilon^2}{2} h_p^\ell \left(R_{ijk}^q h_p^q - R_{ijq}^p h_k^q \right) + \mathcal{O}(\varepsilon^3). \end{aligned}$$

(3) In local coordinates, the Ricci tensor $\widetilde{\text{Rc}}$ of $\tilde{g}$ is

$$\begin{aligned}\tilde{R}_{ij} = R_{ij} - \frac{\varepsilon}{2} & \left[\Delta_\ell h_{ij} + \nabla_i \nabla_j H + \nabla_i (\delta h)_j + \nabla_j (\delta h)_i \right] \\ & + \frac{\varepsilon^2}{2} h^{pq} (\nabla_p \nabla_q h_{ij} + \nabla_i \nabla_j h_{pq} - \nabla_i \nabla_p h_{qj} - \nabla_j \nabla_p h_{qi}) \\ & + \frac{\varepsilon^2}{2} \left(\frac{1}{2} \nabla_i h_p^q \nabla_j h_q^p + \nabla_p h_i^q \nabla^p h_{qj} - \nabla_p h_i^q \nabla_q h_j^p \right) \\ & + \frac{\varepsilon^2}{2} \left(\frac{1}{2} \nabla_k H + (\delta h)_k \right) (\nabla_i h_j^k + \nabla_j h_i^k - \nabla^k h_{ij}) \\ & + \frac{\varepsilon^2}{2} h^{pq} (h_i^k R_{jpkq} + h_j^k R_{ipkq} + 2h_q^k R_{ipkj}) + \mathcal{O}(\varepsilon^3).\end{aligned}$$

(4) If X and Y are vector fields on $\mathcal{M}^m$ and $\tilde{X} = X + \varepsilon Y$, then

$$\mathcal{L}_{\tilde{X}} \tilde{g} = \mathcal{L}_X g + \varepsilon (\mathcal{L}_Y g + \mathcal{L}_X h) + \varepsilon^2 \mathcal{L}_Y h;$$

in particular,

$$\begin{aligned}(\mathcal{L}_{\tilde{X}} \tilde{g})_{ij} &= \nabla_i X_j + \nabla_j X_i \\ &+ \varepsilon [(\nabla_i Y_j + \nabla_j Y_i) + (X^k \nabla_k h_{ij} + \nabla_i X^k h_{kj} + \nabla_j X^k h_{ik})] \\ &+ \varepsilon^2 (Y^k \nabla_k h_{ij} + \nabla_i Y^k h_{kj} + \nabla_j Y^k h_{ik}).\end{aligned}$$

3.4. Arbitrary perturbations of a stationary solution. Let $(\mathcal{M}^m, g, X)$ be any stationary solution of the dilated Ricci flow (72), and let h be an arbitrary symmetric $(2, 0)$ -tensor such that the perturbation

$$(75) \quad \tilde{g} = g + h$$

is a Riemannian metric on $\mathcal{M}^m$. Denote the trace of h with respect to g by

$$(76) \quad H = \text{tr}_g h.$$

Let Y be the vector field metrically dual to the 1-form $\frac{1}{2}dH + \delta h$, namely

$$(77) \quad Y = \left(\frac{1}{2}dH + \delta h \right)^\sharp,$$

and form the perturbation

$$(78) \quad \tilde{X} = X + Y$$

of the vector field X . The role of Y is to implement a DeTurck trick [9, 10] that makes the linearization of the dilated Ricci flow (72) strictly parabolic. (In the elliptic context, this choice is frequently called the Bianchi gauge [6].) Now applying Lemma 28 gives the following result.

Lemma 29. *The metric $\tilde{g}(\tau)$ will be a solution of the dilated Ricci flow (72) for the vector field $\tilde{X}$ if and only if h evolves by the nonlinear system*

$$(79) \quad \frac{\partial}{\partial \tau} h = \mathcal{A}h + \mathcal{Q}(h) + \mathcal{C}(h),$$

where $\mathcal{A}$ is the linear elliptic operator defined by

$$(80) \quad \mathcal{A}h = \Delta_\ell h - \mathcal{L}_X h + h,$$

the quadratic term $\mathcal{Q}(h)$ is defined by

$$\begin{aligned}
(81a) \quad [\mathcal{Q}(h)]_{ij} &= h_i^p \nabla_j \nabla_q h_p^q + h_j^p \nabla_i \nabla_q h_p^q \\
(81b) \quad &+ h^{pq} (\nabla_i \nabla_p h_{qj} + \nabla_j \nabla_p h_{qi} - \nabla_p \nabla_q h_{ij} - \nabla_i \nabla_j h_{pq}) \\
(81c) \quad &- \frac{1}{2} (h_i^k \nabla_j \nabla_k H + h_j^k \nabla_i \nabla_k H + \nabla_i h_j^k \nabla_k H + \nabla_j h_i^k \nabla_k H) \\
(81d) \quad &+ \nabla_i h_j^p \nabla_q h_p^q + \nabla_j h_i^p \nabla_q h_p^q + \nabla_p h_i^q \nabla_q h_j^p \\
(81e) \quad &- \nabla_p h_i^q \nabla^p h_{jq} - \frac{1}{2} \nabla_i h_p^q \nabla_j h_q^p \\
(81f) \quad &+ h^{pq} (h_i^k R_{jpqk} + h_j^k R_{ipqk} - 2h_q^k R_{ipkj}),
\end{aligned}$$

and $\mathcal{C}(h)$ is at least third-order in h and its covariant derivatives.

The cubic term $\mathcal{C}(h)$ will not be important in the formal asymptotic analysis that follows.

3.5. The linearization at the cylinder soliton. In the case that the stationary solution of the dilated Ricci flow is the cylinder soliton $(\mathbb{R} \times S^n, g, X)$ introduced in § 3.1, the linear operator $\mathcal{A}$ defined in (80) can be written in a more useful form. To see this, we again work in the coordinate system $(y^0; y^1, \dots, y^n)$, with $y^0 \equiv x \in \mathbb{R}$ and $(y^1, \dots, y^n) \equiv y \in S^n$. Recalling formula (69), it is easy to see that

$$h_{ij} - (R_i^k h_{kj} + R_j^k h_{ik}) = \begin{cases} h_{00} & \text{if } i = j = 0 \\ h_{i0}/2 & \text{if } 1 \leq i \leq n \\ h_{0j}/2 & \text{if } 1 \leq j \leq n \\ 0 & \text{if } 1 \leq i, j \leq n. \end{cases}$$

This observation allows one to write the Lie derivative of h with respect to X as

$$\begin{aligned}
(\mathcal{L}_X h)_{ij} &= \frac{x}{2} \frac{\partial}{\partial x} h_{ij} + \frac{1}{2} h_{ij} (\delta_i^0 + \delta_j^0) \\
&= \frac{x}{2} \frac{\partial}{\partial x} h_{ij} + h_{ij} - (R_i^k h_{kj} + R_j^k h_{ik}).
\end{aligned}$$

Then using the simple identity

$$R_{i\ell} R_{jk} - R_{ik} R_{j\ell} = \frac{n-1}{2} R_{ijk\ell},$$

one reaches the following conclusion.

Lemma 30. *On the cylinder soliton, the linear operator $\mathcal{A}$ defined in (80) is given by*

$$(82) \quad (\mathcal{A}h)_{ij} = \Delta h_{ij} - \frac{x}{2} \frac{\partial}{\partial x} h_{ij} + \frac{4}{n-1} (R_{ij} R_{pq} - R_{iq} R_{pj}) h^{pq},$$

where Δh denotes the rough Laplacian.

3.6. An upper bound for the spectrum of the linearization. Since the manifold $\mathbb{R} \times S^n$ is noncompact, it is not necessarily the case that the formally elliptic operator $\mathcal{A}$ has discrete point spectrum. For our operator $\mathcal{A}$ the first order term $-\frac{1}{2}x\partial_x$ makes this happen, as we will now show.

We consider the bundle $\mathcal{S}_2$ of $(2, 0)$ -tensors on $\mathbb{R} \times S^n$ and the space $C_0^\infty(\mathcal{S}_2)$ of smooth compactly supported sections of this bundle. The operator $\mathcal{A}$ maps this space to itself. We define the inner product

$$(h, k) \doteq \int_{\mathbb{R}} \int_{S^n} h_{ij} k_{ij} e^{-x^2/4} d\theta dx$$

where $d\theta$ is the volume form on S^n . We write $\|h\| = \sqrt{(h, h)}$ for the corresponding norm. The completion of $C_0^\infty(\mathcal{S}_2)$ with respect to this norm is $L^2(\mathcal{S}_2; e^{-x^2/4} d\theta dx)$, which we abbreviate to L^2 . (See [11], for instance.)

Lemma 31. *The closure of the densely defined operator $\mathcal{A}$ is a self-adjoint operator with compact resolvent. $\mathcal{A}$ is bounded from above. Its spectrum consists of discrete finite multiplicity point spectrum.*

Proof. One can write $\mathcal{A}$ as

$$(83) \quad (\mathcal{A}h)_{ij} = e^{x^2/4} \frac{\partial}{\partial x} \left\{ e^{-x^2/4} \frac{\partial h_{ij}}{\partial x} \right\} + \frac{1}{2(n-1)} \hat{\Delta} h_{ij} + V_{ijpq} h^{pq}$$

in which $\hat{\Delta}$ is the rough Laplacian in the direction tangential to the spheres $\{x\} \times S^n$ and

$$V_{ijpq} \doteq \frac{4}{n-1} (R_{ij} R_{pq} - R_{iq} R_{jp}).$$

By integration by parts, one finds that for any $h, k \in C_0^\infty(\mathcal{S}_2)$ one has

$$(84) \quad (h, -\mathcal{A}k) = \int_{\mathbb{R}} \int_{S^n} \left\{ \partial_x h_{ij} \partial_x k_{ij} + \frac{1}{2(n-1)} \hat{\nabla}_\ell h_{ij} \hat{\nabla}_\ell k_{ij} - V_{ijpq} h_{ij} k_{pq} \right\} e^{-x^2/4} d\theta dx,$$

from which it is evident that $\mathcal{A}$ is symmetric. Boundedness of V_{ijpq} implies that there is a constant C_0 such that

$$(85) \quad (h, -\mathcal{A}h) \geq \int_{\mathbb{R}} \int_{S^n} |\nabla h|^2 e^{-x^2/4} d\theta dx - C_0 \|h\|^2$$

holds for all $h \in C_0^\infty(\mathcal{S}_2)$.

Let $H^1 = H^1(\mathcal{S}_2)$ be the completion of $C_0^\infty(\mathcal{S}_2)$ for the norm corresponding to the inner product

$$\langle h, k \rangle \doteq \int_{\mathbb{R}} \int_{S^n} \left\{ \nabla_i h_{pq} \nabla_i k_{pq} + h_{pq} k_{pq} \right\} e^{-x^2/4} d\theta dx.$$

(Again, see [11].) Then the standard Hilbert space arguments show that for all $\lambda > C_0$, the equation $\lambda h - \mathcal{A}h = k$ has a distributional solution $h \in H^1$ for any $k \in L^2$. It follows that the closure of $\mathcal{A}$ is a self-adjoint operator in L^2 whose domain

$$D(\mathcal{A}) = \{(\lambda - \mathcal{A})^{-1}k \mid k \in L^2\}$$

is contained in H^1 . (Here it makes no difference which $\lambda > C_0$ one chooses.) In fact, the inequality (85) implies that the H^1 norm is equivalent with the norm $\|h\| = \sqrt{(h, (\lambda - \mathcal{A})h)}$, so that H^1 is the form domain of $\mathcal{A}$, i.e. $H^1 = D(\sqrt{\lambda - \mathcal{A}})$

The inequality (85) tells us that the spectrum of $\mathcal{A}$ is contained in the interval $(-\infty, C_0]$.

To see that $\mathcal{A}$ has compact resolvent, we recall Lemma 10, which implies

$$\|xh\|^2 \leq C \int_{\mathbb{R}} \int_{S^n} \{|\partial_x h|^2 + |h|^2\} e^{-x^2/4} d\theta dx \leq C \|h\|_{H^1}^2.$$

Hence

$$\|h\|^2 + \|xh\|^2 + \|\nabla h\|^2 \leq C(h, (\lambda - \mathcal{A})h).$$

This inequality together with the Rellich-Kondrachov theorem imply that the imbedding $H^1 \subset L^2$ is compact. Since H^1 is the form domain of $\mathcal{A}$, we conclude that $\mathcal{A}$ indeed has compact resolvent, so that its spectrum is pure point spectrum. $\square$

REMARK. More generally, the linearization of the modified Ricci flow operator at any gradient Ricci soliton $(\mathcal{M}^n, g)$ will be self-adjoint in $L^2(\mathcal{M}^n, g; e^{-\varphi} d\mu)$ if $d\mu$ is the measure naturally associated to g and one takes φ to be the soliton potential function.

3.7. A decomposition into invariant subspaces. In order to compute the spectrum of $\mathcal{A}$, we split the space $L^2(\mathcal{S}_2)$ into a number of $\mathcal{A}$ -invariant subspaces.

It is a standard fact that any $(2, 0)$ -tensor h on a Riemannian manifold $(\mathcal{M}^m, g)$ admits the decomposition

$$h = h^\vee + h^\circ,$$

where

$$h^\vee = (\text{tr}_g h) g$$

is a multiple of the metric, and

$$h^\circ = h - \frac{1}{m} h^\vee$$

is trace free.

One can exploit the product structure of the cylinder soliton $(\mathbb{R} \times S^n, g, X)$ to obtain a more refined decomposition. Again using coordinates $(y^0; y^1, \dots, y^n)$ and adopting the convention that Roman indices lie in the range $0 \dots n$ while Greek indices lie in $1 \dots n$, we decompose a given symmetric $(2, 0)$ -tensor h on $\mathbb{R} \times S^n$ as

$$(86a) \quad h = M dx \otimes dx + N_\alpha (dx \otimes dy^\alpha + dy^\alpha \otimes dx) + (O_{\alpha\beta}^\vee + O_{\alpha\beta}^\circ) dy^\alpha \otimes dy^\beta$$

$$(86b) \quad = M dx \otimes dx + N_\alpha (dx \otimes dy^\alpha + dy^\alpha \otimes dx) + (P \hat{g}_{\alpha\beta} + Q_{\alpha\beta}) dy^\alpha \otimes dy^\beta.$$

Here one should regard M as a scalar function on $\mathbb{R} \times S^n$, $N(x, \cdot)$ as a one-parameter family of 1-forms defined on the spheres $\{x\} \times S^n$, P as the scalar function

$$P = \frac{2(n-1)}{n} (H - M),$$

and $Q(x, \cdot)$ as the one-parameter family of trace-free $(2, 0)$ -tensors defined on the spheres $\{x\} \times S^n$ by

$$Q_{\alpha\beta} = h_{\alpha\beta} - P \hat{g}_{\alpha\beta}.$$

The utility of this decomposition is that it exhibits a set of invariant subspaces for $\mathcal{A}$. To demonstrate this, it will be helpful to fix additional notation. Let $\hat{\Delta}$

denote the rough Laplacian of the canonical sphere metric $\hat{g}$. Let $-\hat{\Delta}_{dR}$ denote its Hodge-de Rham Laplacian, recalling that $\hat{\Delta}_{dR}$ acts on a 1-form N by

$$\hat{\Delta}_{dR} N_\alpha = -[(d\delta + \delta d) N]_\alpha = \hat{\Delta} N_\alpha - \hat{R}_\alpha^\beta N_\beta = \hat{\Delta} N_\alpha - (n-1) N_\alpha.$$

Let $\hat{\Delta}_\ell$ denote the Lichnerowicz Laplacian of $\hat{g}$, which acts on a trace-free tensor Q by

$$\hat{\Delta}_\ell Q_{\alpha\beta} = \hat{\Delta} Q_{\alpha\beta} - 2n Q_{\alpha\beta}.$$

Finally, let $\mathcal{B}$ denote the differential operator defined by

$$(87) \quad \mathcal{B} = \frac{\partial^2}{\partial x^2} - \frac{x}{2} \frac{\partial}{\partial x} = e^{x^2/4} \frac{\partial}{\partial x} e^{-x^2/4} \frac{\partial}{\partial x}.$$

Using these conventions, one computes using equation (82) that the decomposition of $\mathcal{A}h$ corresponding to (86) is

$$\begin{aligned} (\mathcal{A}H)_{00} &= \left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} \right] M \\ (\mathcal{A}H)_{\alpha 0} &= \left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} \right] N_\alpha = \left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta}_{dR} + \frac{1}{2} \right] N_\alpha \\ (\mathcal{A}H)_{\alpha\beta}^\vee &= \left\{ \left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} + 1 \right] P \right\} \hat{g}_{\alpha\beta} \\ (\mathcal{A}H)_{\alpha\beta}^\circ &= \left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} - \frac{1}{n-1} \right] Q_{\alpha\beta}. \end{aligned}$$

This calculation proves the following result.

Lemma 32. *On the cylinder soliton, the operator $\mathcal{A}$ given by (82) may be represented schematically with respect to the decomposition (86) as*

$$(88) \quad \mathcal{A} : \begin{pmatrix} M \\ N \\ P \\ Q \end{pmatrix} \mapsto \begin{pmatrix} \left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} \right] M \\ \left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta}_{dR} + \frac{1}{2} \right] N \\ \left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} + 1 \right] P \\ \left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} - \frac{1}{n-1} \right] Q \end{pmatrix}.$$

The only contribution to the spectrum of $\mathcal{A}$ comes from its four component operators: $\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta}$ and $\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} + 1$ acting on scalar functions, $\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta}_{dR} + \frac{1}{2}$ acting on 1-forms, and $\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} - \frac{1}{n-1}$ acting on trace-free $(2, 0)$ -tensors.

3.8. The spectrum of the linearization. The decomposition obtained in Lemma 32 allows us to analyze the spectrum of $\mathcal{A}$ using separation of variables. To illustrate the idea, suppose that

$$M(x, y) = h(x) F(y),$$

where h is an eigenfunction of $\mathcal{B}$ with eigenvalue λ , and F is an eigenfunction of $\frac{1}{2(n-1)} \hat{\Delta}$ with eigenvalue μ . Then

$$\left[\mathcal{B} + \frac{1}{2(n-1)} \hat{\Delta} \right] M = (\lambda + \mu) M.$$

This line of argument leads easily to the following observation.

Lemma 33. *All eigenvalues of $\mathcal{A} : D(\mathcal{A}) \rightarrow L^2$ have the form $\lambda + \mu$, where λ is an eigenvalue of $\mathcal{B}$ acting on $\mathfrak{h} = L^2(\mathbb{R}; e^{-x^2/4})$, and μ is an eigenvalue of either $\frac{1}{2(n-1)}\hat{\Delta}$ or $\frac{1}{2(n-1)}\hat{\Delta} + 1$ acting on $C^\infty(S^n)$, of $\frac{1}{2(n-1)}\hat{\Delta}_{dR} + \frac{1}{2}$ acting on $\Omega_1(S^n)$, or of $\frac{1}{2(n-1)}\hat{\Delta} - \frac{1}{n-1}$ acting on the space of smooth trace-free $(2,0)$ -tensors on S^n .*

To make this result useful, we need some simple facts about the component operators that appear in (88).

THE OPERATOR $\mathcal{B} = \frac{\partial^2}{\partial x^2} - \frac{x}{2} \frac{\partial}{\partial x}$ ACTING ON SCALAR FUNCTIONS. The Hermite polynomials $\{h_k\}_{k=0}^\infty$ constitute a complete orthogonal family for the operator $\mathcal{B}$ on the weighted Hilbert space

$$\mathfrak{h} = L^2\left(\mathbb{R}, e^{-x^2/4} dx\right).$$

Moreover, one has

$$(\mathcal{B} + 1)h_k = \left(1 - \frac{k}{2}\right)h_k,$$

so that the spectrum of $\mathcal{B}$ is $\{0, -\frac{1}{2}, -1, -\frac{3}{2}, \dots\}$. (Notice that $\mathfrak{h}$ is a larger space than we defined in Section 2, since we not restricting to even functions here.)

THE OPERATOR $\frac{1}{2(n-1)}\hat{\Delta}$ ACTING ON SCALAR FUNCTIONS. It is well known that the spectrum of $\hat{\Delta}$ acting on $C^\infty(S^n)$ is $\{\lambda_k\}_{k \geq 0}$, where

$$\lambda_k = -k(n + k - 1).$$

In particular, the only nonnegative eigenvalue of $\frac{1}{2(n-1)}\hat{\Delta}$ is $\lambda_0 = 0$. Its eigenspace consists of the constant functions.

THE OPERATOR $\frac{1}{2(n-1)}\hat{\Delta} + 1$ ACTING ON SCALAR FUNCTIONS. If $k \geq 2$, then

$$\lambda_k + 2(n-1) = (2-k)(n-1) - k^2 \leq -k^2 < 0.$$

Hence the only possible nonnegative eigenvalues of $\frac{1}{2(n-1)}\hat{\Delta} + 1$ are

$$\frac{\lambda_0 + 2(n-1)}{2(n-1)} = 1$$

and

$$\frac{\lambda_1 + 2(n-1)}{2(n-1)} = \frac{1}{2} \frac{n-2}{n-1} \in \left[0, \frac{1}{2}\right).$$

The eigenspace corresponding to 1 consists of constants. The eigenspace corresponding to $\frac{1}{2} \frac{n-2}{n-1}$ consists of the spherical harmonics: the restrictions to S^n of the linear functions on $\mathbb{R}^{n+1}$.

THE OPERATOR $\frac{1}{2(n-1)}\hat{\Delta}_{dR} + \frac{1}{2}$ ACTING ON 1-FORMS. It is shown in [7] that the spectrum of $\hat{\Delta}_{dR}$ acting on the space $\Omega_1(S^n)$ of smooth 1-forms is

$$\{\lambda_k\}_{k \geq 1} \cup \{\lambda_k + 2 - n\}_{k \geq 1}.$$

If $k \geq 1$, then $\lambda_k + n - 1 \leq -1$. Because $\lambda_k + 2 - n \leq \lambda_k$ in all dimensions $n \geq 2$, it follows that $\frac{1}{2(n-1)}\hat{\Delta}_{dR} + \frac{1}{2}$ is negative definite.

THE OPERATOR $\frac{1}{2(n-1)}\hat{\Delta} - \frac{1}{n-1}$ ACTING ON TRACE-FREE $(2,0)$ -TENSORS. The operator $\frac{1}{2(n-1)}\hat{\Delta} - \frac{1}{n-1}$ is clearly negative definite.

Combining these observations with Lemma 33 leads to the following conclusion.

Lemma 34. *On the cylinder soliton, the only possible non-negative eigenvalues of the linearization $\mathcal{A}$ are 0, $\frac{1}{2}\frac{n-2}{n-1}$, $\frac{1}{2}$, and 1.*

- The eigenspace corresponding to 1 consists of constant multiples of $\hat{g}$.
- The eigenspace corresponding to $\frac{1}{2}$ consists of multiples of $\hat{g}$ that are linear in $x \in \mathbb{R}$.
- The eigenspace corresponding to $\frac{1}{2}\frac{n-2}{n-1}$ consists of spherical harmonics.
- The eigenspace corresponding to 0 consists of constant functions and of multiples of $\hat{g}$ that are quadratic in $x \in \mathbb{R}$. (When $n = 2$, the spherical harmonics also belong to this eigenspace.)

REMARK. The results above can be obtained in another way. Since all geometric data of the metric (64) are independent of $x \in \mathbb{R}$, one has the simple commutator

$$\left[\mathcal{A}, \frac{\partial}{\partial x} \right] = \frac{1}{2} \frac{\partial}{\partial x}.$$

So if $\mathcal{A}h = \lambda h$, a straightforward induction argument shows that

$$\mathcal{A} \left(\frac{\partial^k}{\partial x^k} h \right) = \left(\lambda + \frac{k}{2} \right) \left(\frac{\partial^k}{\partial x^k} h \right)$$

for all integers $k \geq 0$. Since Lemma 31 proves that the spectrum of $\mathcal{A}$ is bounded from above, it follows readily that for every eigentensor h of $\mathcal{A}$, there exist some integer $\kappa = \kappa(h) \geq 0$ and a family of $(2,0)$ -tensors $\hat{h}_1, \dots, \hat{h}_\kappa$ independent of $x \in \mathbb{R}$ such that

$$h = \sum_{k=0}^{\kappa} x^k \hat{h}_k.$$

3.9. Our Ansatz. The analysis in § 3.8 shows that any eigenvalues of $\mathcal{A}$ corresponding to trace-free (i.e. non-rotationally symmetric) perturbations of the cylinder soliton are all strictly negative. As we explained in the introduction to this section, the only perturbations relevant to our formal center-manifold computation are those corresponding to null eigenvalues of the linearization. Therefore, we adopt the *Ansatz* that

$$h = M dx \otimes dx + N_\alpha (dx \otimes dy^\alpha + dy^\alpha \otimes dx) + (P\hat{g}_{\alpha\beta} + Q_{\alpha\beta}) dy^\alpha \otimes dy^\beta$$

has the special form

$$(89) \quad h(x, y, \tau) = u(x, \tau) dx \otimes dx + \{v(x, \tau) + w(y, \tau)\} \hat{g},$$

where once again, τ denotes rescaled time, x denotes a point in $\mathbb{R}$, and y denotes a point in S^n . Specifically — continuing to let $\{h_k\}_{k=0}^\infty$ denote the Hermite polynomials — we assume that $M \equiv u$, where

$$(90) \quad u(x, \tau) = u_0(\tau) h_0(x) = u_0(\tau).$$

We further assume that $N \equiv 0$, that $Q \equiv 0$, and that $P \equiv v + w$, where

$$(91) \quad v(x, \tau) = v_2(\tau) h_2(x) = v_2(\tau) (x^2 - 2)$$

and

$$(92) \quad w(y, \tau) = \begin{cases} \sum_{i=1}^3 \omega_i(\tau) \Omega_i(y) & \text{if } n = 2 \\ 0 & \text{otherwise.} \end{cases}$$

Here $(\Omega_1, \Omega_2, \Omega_3)$ is a basis for the space of spherical harmonics on S^2 . (Recall that w corresponds to a positive eigenvalue of the linearization whenever the *total dimension* $n + 1$ is at least 4.)

To implement our formal center-manifold analysis, we shall study the flow

$$(93) \quad \frac{\partial}{\partial \tau} h = \mathcal{A}h + \mathcal{Q}(h)$$

that models equation (79) up to second order. To do so requires us to analyze the quadratic term $\mathcal{Q}(h)$ at the cylinder soliton in a manner analogous to what was done above for the linearization $\mathcal{A}$. To simplify the notation, we again use coordinates $(y^0; y^1, \dots, y^n)$ and assume that Roman indices lie in the range $0 \dots n$ while Greek indices lie in $1 \dots n$. If f is a smooth function, we further adopt the convention that $f_x = \partial f / \partial x$ and $f_\alpha = \partial f / \partial y^\alpha$.

We shall begin with computations that are more general than what we need at the moment. To wit, we assume only that h has the form given by (89), without imposing the specific assumptions (90)–(92). One then verifies readily that all first covariant derivatives of the tensor h vanish except

$$\begin{aligned} \nabla_0 h_{00} &= u_x \\ \nabla_0 h_{\alpha\beta} &= v_x \hat{g}_{\alpha\beta} \\ \nabla_\gamma h_{\alpha\beta} &= w_\gamma \hat{g}_{\alpha\beta}. \end{aligned}$$

All second covariant derivatives of h vanish except

$$\begin{aligned} \nabla_0 \nabla_0 h_{00} &= u_{xx} \\ \nabla_0 \nabla_0 h_{\alpha\beta} &= v_{xx} \hat{g}_{\alpha\beta} \\ \nabla_\gamma \nabla_\delta h_{\alpha\beta} &= \hat{\nabla}_\gamma \hat{\nabla}_\delta w \hat{g}_{\alpha\beta}. \end{aligned}$$

Moreover, the only nonvanishing derivatives of the trace

$$H = g^{ij} h_{ij} = u + \frac{n}{2(n-1)}(v + w)$$

are

$$\begin{aligned} \nabla_0 H &= u_x + \frac{n}{2(n-1)}v_x \\ \nabla_\gamma H &= \frac{n}{2(n-1)}w_\gamma \end{aligned}$$

and

$$\begin{aligned} \nabla_0 \nabla_0 H &= u_{xx} + \frac{n}{2(n-1)}v_{xx} \\ \nabla_\gamma \nabla_\delta H &= \frac{n}{2(n-1)}\hat{\nabla}_\gamma \hat{\nabla}_\delta w. \end{aligned}$$

Using these formulas, a straightforward calculation reveals that the first component of the quadratic term

$$\begin{aligned} [\mathcal{Q}(h)]_{00} = & 2h_{00}\nabla_0\nabla_0h_{00} - h^{\alpha\beta}\nabla_0\nabla_0h_{\alpha\beta} - h_{00}\nabla_0\nabla_0H \\ & - \nabla_0h_{00}\nabla_0H + \frac{3}{2}(\nabla_0h_{00})^2 - \frac{1}{2}\nabla_0h_\alpha^\beta\nabla_0h_\beta^\alpha \end{aligned}$$

may be written as

$$(94a) \quad [\mathcal{Q}(h)]_{00} = uu_{xx} - \frac{n}{4(n-1)^2}v_{xx}[2(n-1)u + v + w]$$

$$(94b) \quad + \frac{1}{2}u_x^2 - \frac{n}{8(n-1)^2}v_x[4(n-1)u_x + v_x].$$

Then using the fact that

$$g^{\gamma\delta}\hat{\nabla}_\gamma\hat{\nabla}_\delta w = \frac{1}{2(n-1)}\hat{\Delta}w = -\frac{n}{2(n-1)}w,$$

one computes that the remaining components

$$\begin{aligned} [\mathcal{Q}(h)]_{\alpha\beta} = & h_\alpha^\gamma\nabla_\beta\nabla_\delta h_\gamma^\delta + h_\beta^\gamma\nabla_\alpha\nabla_\delta h_\gamma^\delta - h^{00}\nabla_0\nabla_0h_{\alpha\beta} \\ & + h^{\gamma\delta}(\nabla_\alpha\nabla_\gamma h_{\delta\beta} + \nabla_\beta\nabla_\gamma h_{\delta\alpha} - \nabla_\gamma\nabla_\delta h_{\alpha\beta} - \nabla_\alpha\nabla_\beta h_{\gamma\delta}) \\ & - \frac{1}{2}\left(h_\alpha^\gamma\nabla_\beta\nabla_\gamma H + h_\beta^\gamma\nabla_\alpha\nabla_\gamma H + \nabla_\alpha h_\beta^\gamma\nabla_\gamma H + \nabla_\beta h_\alpha^\gamma\nabla_\gamma H\right) \\ & + \nabla_\alpha h_\beta^\gamma\nabla_\delta h_\gamma^\delta + \nabla_\beta h_\alpha^\gamma\nabla_\delta h_\gamma^\delta + \nabla_\gamma h_\alpha^\delta\nabla_\delta h_\beta^\gamma \\ & - \nabla_0 h_\alpha^\gamma\nabla_0 h_{\beta\gamma} - \nabla_\delta h_\alpha^\gamma\nabla^\delta h_{\beta\gamma} - \frac{1}{2}\nabla_\alpha h_\gamma^\delta\nabla_\beta h_\delta^\gamma \\ & + h^{\gamma\delta}\left(h_\alpha^\eta R_{\beta\gamma\delta\eta} + h_\beta^\eta R_{\alpha\gamma\delta\eta} - 2h_\delta^\eta R_{\alpha\gamma\eta\beta}\right) \end{aligned}$$

are given by

$$(95a) \quad [\mathcal{Q}(h)]_{\alpha\beta} = -\frac{1}{2(n-1)}\{2(n-1)uv_{xx} + v_x^2\}\hat{g}_{\alpha\beta}$$

$$(95b) \quad -\frac{1}{4(n-1)^2}\left\{\left|\hat{\nabla}w\right|_{\hat{g}}^2 - nw(v+w)\right\}\hat{g}_{\alpha\beta}$$

$$(95c) \quad -\frac{n-2}{8(n-1)^2}\left\{4(v+w)\hat{\nabla}_\alpha\hat{\nabla}_\beta w + 3w_\alpha w_\beta\right\}.$$

3.10. The inner layer $|x| = o(\sqrt{\tau})$. We now derive a formal profile of a solution

$$\tilde{g}(x, y, \tau) = g(x, y) + h(x, y, \tau)$$

near a singularity modeled on the cylinder soliton g . We assume that (89) and (90)–(92) hold. These assumptions force h to belong to the kernel of the linearization of equation (79), because

$$\mathcal{A}h = [\mathcal{B}u]dx \otimes dx + [(\mathcal{B} + 1)v]\hat{g} = 0.$$

Recall that w vanishes by assumption if $n > 2$. In this context, (94) becomes

$$(96) \quad [\mathcal{Q}(h)]_{00} = -\left\{v_2w + \frac{n}{(n-1)^2}[(n-1)u_0v_2 + v_2^2]\right\}h_0 - \left\{\frac{n}{(n-1)^2}v_2^2\right\}h_2.$$

And (95) reduces to

$$[\mathcal{Q}(h)]_{\alpha\beta} = \mathcal{P}\hat{g}_{\alpha\beta},$$

where

$$(97) \quad \mathcal{P} = \left\{ \frac{1}{2} w^2 - \frac{1}{4} \left| \hat{\nabla} w \right|_{\hat{g}}^2 - 2u_0 v_2 - \frac{4}{n-1} v_2^2 \right\} h_0 + \left\{ \frac{1}{2} v_2 w - \frac{2}{n-1} v_2^2 \right\} h_2.$$

In order efficiently to derive ODE for the functions u_0 , v_2 , and $\omega_1, \omega_2, \omega_3$ from equations (96) and (97), it is helpful to adopt some additional notation: if f is a function of $y \in S^n$, we define

$$\|f\|_{S^n}^2 = \int_{S^n} f(y)^2 d\hat{\mu};$$

and if X and Y are symmetric $(2, 0)$ -tensors on $\mathbb{R} \times S^n$, we define

$$\langle\langle X, Y \rangle\rangle = \int_{-\infty}^{\infty} \langle X, Y \rangle_x e^{-x^2/4} dx,$$

where $\langle X, Y \rangle_x = \int_{\{x\} \times S^n} \langle X, Y \rangle_{\hat{g}} d\hat{\mu}$. By using the fact that $\{h_k\}_{k=0}^{\infty}$ is an orthogonal family for $\mathfrak{h}$, it is then easy to see that

$$\begin{aligned} u'_0(\tau) \cdot \|h_0\|_{\mathfrak{h}}^2 \|1\|_{S^n}^2 &= \frac{d}{dt} \langle\langle h, h_0 dx^2 \rangle\rangle \\ &= \langle\langle \mathcal{Q}(h), h_0 dx^2 \rangle\rangle \\ &= -\frac{n}{(n-1)^2} [(n-1)u_0 v_2 + v_2^2] \cdot \|h_0\|_{\mathfrak{h}}^2 \|1\|_{S^n}^2 \end{aligned}$$

and

$$\begin{aligned} v'_2(\tau) \cdot n \|h_2\|_{\mathfrak{h}}^2 \|1\|_{S^n}^2 &= \frac{d}{dt} \langle\langle h, h_2 \hat{g} \rangle\rangle \\ &= \langle\langle \mathcal{Q}(h), h_2 \hat{g} \rangle\rangle \\ &= -\frac{2}{n-1} v_2^2 \cdot n \|h_2\|_{\mathfrak{h}}^2 \|1\|_{S^n}^2. \end{aligned}$$

Similarly, for any $i \in \{1, 2, 3\}$, we find when $n = 2$ that

$$\omega'_i(\tau) \cdot 2 \|h_0\|_{\mathfrak{h}}^2 \|\Omega_i\|_{S^2}^2 = \frac{d}{dt} \langle\langle h, \Omega_i \hat{g} \rangle\rangle = \langle\langle \mathcal{Q}(h), \Omega_i \hat{g} \rangle\rangle = 0.$$

This calculations let us derive a formal profile of h in the inner layer $|x| = o(\sqrt{\tau})$.

Lemma 35. *For perturbations of the form*

$$h = \{u_0(\tau) h_0(x)\} dx \otimes dx + \left\{ v_2(\tau) h_2(x) + \sum_{i=1}^3 \omega_i(\tau) \Omega_i(y) \right\} \hat{g},$$

the flow (93) is equivalent to the system of ODE

$$(98) \quad \frac{d}{d\tau} u_0 = -\frac{n}{(n-1)^2} [(n-1)u_0 v_2 + v_2^2]$$

$$(99) \quad \frac{d}{d\tau} v_2 = -\frac{2}{n-1} v_2^2$$

$$(100) \quad \frac{d}{dt} \omega_i = 0 \quad (i = 1, 2, 3)$$

whose solutions, up to an error term of $o(\tau^{-1})$ as $\tau \rightarrow \infty$, are

$$(101) \quad u_0(\tau) = \begin{cases} -\frac{\log(2\tau)}{2\tau} + \frac{c}{\tau} & \text{if } n = 2 \\ -\frac{n}{2(n-2)} \frac{1}{\tau} & \text{otherwise} \end{cases}$$

$$(102) \quad v_2(\tau) = \frac{n-1}{2\tau}$$

$$(103) \quad \omega_i(\tau) = \omega_i(0) \quad (i = 1, 2, 3).$$

Formally, this is the profile of h in the region $|x| = o(\sqrt{\tau})$.

Proof. If v_2 is not identically zero, equation (99) has the explicit solution

$$v_2(\tau) = \frac{n-1}{c+2\tau}.$$

Then equation (98) becomes

$$\frac{d}{d\tau} u_0(\tau) + \left(\frac{n}{c+2\tau} \right) u_0 = -\frac{n}{(c+2\tau)^2}.$$

An integrating factor for this linear ODE is $(c+2\tau)^{n/2}$, whence one obtains

$$u_0(\tau) = \begin{cases} -\frac{\log(c+2\tau)}{c+2\tau} + \frac{C}{c+2\tau} & \text{if } n = 2 \\ -\frac{n}{n-2} \frac{1}{c+2\tau} + \frac{C}{(c+2\tau)^{n/2}} & \text{otherwise.} \end{cases}$$

□

REMARK. To compare the formal results above with those obtained rigorously in Section 2, observe that our choices of dilating factors imply that the quantity $a_2(\tau)$ appearing in Section 2 and the quantity $v_2(\tau)$ appearing above are related by

$$a_2(\tau) \approx \frac{v_2(\tau)}{4(n-1)} \approx \frac{1}{8\tau},$$

because in the inner layer, we have $x \approx \sigma$ and

$$\begin{aligned} & \{2(n-1) + v_2(\tau) h_2(x) + o(\tau^{-1})\} \hat{g} \\ &= 2(n-1) [1 + a_2(\tau) h_2(\sigma) + o(\tau^{-1})]^2 \hat{g} \\ &= \{2(n-1) + 4(n-1) a_2(\tau) h_2(\sigma) + o(\tau^{-1})\} \hat{g}. \end{aligned}$$

REMARK. By modifying $\tilde{g}$ by an initial conformal diffeomorphism, one may make all ω_i vanish. Therefore, throughout the remainder of this paper, we will assume that $\omega_i \equiv 0$ for $i = 1, 2, 3$.

3.11. The intermediate layer $|x| = \mathcal{O}(\sqrt{\tau})$. In order to study the region where $|x| = \mathcal{O}(\sqrt{\tau})$, we replace the coordinates (x, τ) with (ξ, τ) , where

$$(104) \quad \xi = \frac{x}{\sqrt{\tau}}.$$

If f is a smooth function of (x, τ) , we define $F(\xi, \tau) = f(x, \tau)$. Then one has the formulas

$$f_\tau = F_\tau - \frac{\xi}{2\tau} F_\xi \quad \text{and} \quad f_x = \frac{1}{\sqrt{\tau}} F_\xi.$$

Let

$$U(\xi, \tau) = u(x, \tau) \quad \text{and} \quad V(\xi, \tau) = v(x, \tau),$$

where u and v are the quantities that appear in the simple *Ansatz* (89). Then substituting formulas (94) and (95) into the PDE (93), one computes that

$$U_\tau = -\frac{\xi}{2} U_\xi + \frac{1}{\tau} \left\{ \begin{array}{l} (1+U) U_{\xi\xi} + \frac{\xi}{2} U_\xi + \frac{1}{2} U_\xi^2 \\ -\frac{n}{4(n-1)^2} V_{\xi\xi} [2(n-1)U + V] \\ -\frac{n}{8(n-1)^2} V_\xi [4(n-1)U_\xi + V_\xi] \end{array} \right\}$$

and

$$V_\tau = V - \frac{\xi}{2} V_\xi + \frac{1}{\tau} \left\{ V_{\xi\xi} (1-U) + \frac{\xi}{2} V_\xi - \frac{1}{2(n-1)} V_\xi^2 \right\}.$$

Recalling Lemma 35, we will construct an approximate solution by considering the equations

$$U_\tau = -\frac{\xi}{2} U_\xi + o(\tau^{-1})$$

and

$$V_\tau = V - \frac{\xi}{2} V_\xi + \mathcal{O}(\tau^{-2}).$$

These equations suggest that the solution is modeled by the time-independent profiles $U(\xi) \equiv C_1$ and $V(\xi) = C_2 \xi^2$, respectively. Matching this layer with the results of Lemma 35 determines C_1 and C_2 , yielding the following result.

Lemma 36. *In the intermediate layer $|x/\sqrt{\tau}| = |\xi| = \mathcal{O}(1)$, the formal solution is modeled by*

$$u(x, \tau) \approx U\left(\frac{x}{\sqrt{\tau}}\right) \equiv 0 \quad \text{and} \quad v(x, \tau) \approx V\left(\frac{x}{\sqrt{\tau}}\right) = \frac{n-1}{2} \frac{x^2}{\tau}.$$

To wit,

$$\tilde{g} \approx dx \otimes dx + 2(n-1) \left[1 + (\xi/2)^2 \right] \hat{g}.$$

3.12. The outer layer $|x| = o(1/\sqrt{T-t})$. To describe the outer layer, we first introduce the “blown-down” coordinate

$$(105) \quad s = e^{-\tau/2} x = \sqrt{T-t} \cdot x,$$

which should be compared to the metric distance s defined in Section 2.

If one were to follow the method above by defining $\bar{U}(s, \tau) = u(x, \tau)$ and $\bar{V}(s, \tau) = v(x, \tau)$, an easy computation would show that

$$\bar{U}_\tau = \mathcal{O}(T-t) \quad \text{and} \quad \bar{V}_\tau = V + \mathcal{O}(T-t).$$

Unfortunately, the steady-state solutions to these equations fail to convey enough useful information.

Instead, we proceed as follows. Fix a large number $\Xi > 0$. For each s , let t_s be the time such that $|\xi| = \Xi$ at t_s , where ξ is defined by (104). Let $\tau_s = \log \frac{1}{1-t_s}$. Then

$$\Xi = |\xi| = \frac{|x|}{\sqrt{\tau_s}} = \frac{|s|}{\sqrt{(T-t_s)\tau_s}}.$$

This shows that

$$\tau_s = \log(\Xi^2) + \log \frac{1}{s^2} + \log \tau_s,$$

hence that

$$(106) \quad \tau_s = [2 + o(1)] \log \frac{1}{|s|} \quad \text{as} \quad |s| \rightarrow 0.$$

Now we “blow down” the solution of the dilated (“blown-up”) Ricci flow by defining

$$\underline{u}(s, \tau) = e^{-\tau} u(x, \tau) \quad \text{and} \quad \underline{v}(s, \tau) = e^{-\tau} v(x, \tau).$$

Then at time t_s , we compute that

$$\underline{v}(s, \tau_s) = e^{-\tau_s} V\left(\frac{x}{\sqrt{\tau_s}}\right) = \frac{n-1}{2} \frac{s^2}{\tau_s}.$$

When $|s|$ is small and Ξ is large, the undilated solution $\underline{v}$ cannot change much in the short time $t_s \leq t < T$. Hence (106) implies that as $|s| \rightarrow 0$, one has

$$\begin{aligned} \underline{v}(s, \tau_s) &= \frac{n-1}{2} \frac{s^2}{[2 + o(1)] \log \frac{1}{|s|}} [1 + o(1)] \\ &= \frac{n-1}{2} \frac{s^2}{\log \frac{1}{s^2}} [1 + o(1)]. \end{aligned}$$

Since $\underline{u}(s, \tau) = e^{-\tau} U(x/\sqrt{\tau}) \equiv 0$, we have obtained the following result.

Lemma 37. *In the outer layer $|x| = o\left(\frac{1}{\sqrt{T-t}}\right)$, the formal solution is modeled by*

$$u(x, \tau) \approx \frac{1}{T-t} \underline{u}(s, \tau) \quad \text{and} \quad v(x, \tau) \approx \frac{1}{T-t} \underline{v}(s, \tau),$$

where

$$\underline{u}(s, \tau) \equiv 0 \quad \text{and} \quad \underline{v}(s, \tau) = \frac{n-1}{2} \frac{s^2}{\log \frac{1}{s^2}} [1 + o(1)].$$

To wit,

$$v(x, \tau) \approx \frac{n-1}{2} \frac{x^2}{\tau + \log \frac{1}{x^2}}.$$

REFERENCES

- [1] **Altschuler, Steven; Angenent, Sigurd B.; Giga, Yoshikazu.** Mean curvature flow through singularities for surfaces of rotation. *J. Geom. Anal.* **5** (1995), no. 3, 293–358.
- [2] **Angenent, S. B.; Aronson, D. G.** Non-axial self-similar hole filling for the porous medium equation. *J. Amer. Math. Soc.* **14** (2001), no. 4, 737–782 (electronic).
- [3] **Angenent, Sigurd B.; Knopf, Dan.** An example of neckpinching for Ricci flow on S^{n+1} . *Math. Res. Lett.* **11** (2004), no. 4, 493–518.
- [4] **Angenent, S. B.; Velázquez, J. J. L.** Asymptotic shape of cusp singularities in curve shortening. *Duke Math. J.* **77** (1995), no. 1, 71–110.

- [5] **Angenent, Sigurd B.; Velázquez, J. J. L.** Degenerate neckpinches in mean curvature flow. *J. Reine Angew. Math.* **482** (1997), 15–66.
- [6] **Biquard, Olivier.** Métriques d'Einstein asymptotiquement symétriques. *Astérisque* No. 265 (2000).
- [7] **Boucetta, Mohamed.** Spectre des laplaciens de Lichnerowicz sur les sphères et les projectifs réels. (French) [Spectra of Lichnerowicz Laplacians on spheres and real projective spaces] *Publ. Mat.* **43** (1999), no. 2, 451–483.
- [8] **Colding, Tobias H.; Kleiner, Bruce.** Singularity structure in mean curvature flow of mean convex sets. Preprint. <http://www.math.lsa.umich.edu/~bkleiner/research.html>
- [9] **DeTurck, Dennis M.** Deforming metrics in the direction of their Ricci tensors. *J. Differential Geom.* **18** (1983), no. 1, 157–162.
- [10] **DeTurck, Dennis M.** Deforming metrics in the direction of their Ricci tensors, improved version. *Collected Papers on Ricci Flow*. Edited by H.-D. Cao, B. Chow, S.-C. Chu, and S.-T. Yau. Internat. Press, Somerville, MA, 2003.
- [11] **Eichhorn, Jürgen; Fricke, Jan.** The module structure theorem for Sobolev spaces on open manifolds. *Math. Nachr.* **194** (1998), 35–47.
- [12] **Feldman, Mikhail; Ilmanen, Tom; Knopf, Dan.** Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons. *J. Differential Geom.* **65** (2003), no. 2, 169–209.
- [13] **Filippas, Stathis; Kohn, Robert V.** Refined asymptotics for the blowup of $u_t - \Delta u = u^p$. *Comm. Pure Appl. Math.* **45** (1992), no. 7, 821–869.
- [14] **Friedman, Avner; McLeod, Bryce.** Blow-up of positive solutions of semilinear heat equations. *Indiana Univ. Math. J.* **34** (1985), no. 2, 425–447.
- [15] **Giga, Yoshikazu; Kohn, Robert V.** Asymptotically self-similar blow-up of semilinear heat equations. *Comm. Pure Appl. Math.* **38** (1985), no. 3, 297–319.
- [16] **Giga, Yoshikazu; Kohn, Robert V.** Characterizing blowup using similarity variables. *Indiana Univ. Math. J.* **36** (1987), no. 1, 1–40.
- [17] **Giga, Yoshikazu; Kohn, Robert V.** Nondegeneracy of blowup for semilinear heat equations. *Comm. Pure Appl. Math.* **42** (1989), no. 6, 845–884.
- [18] **Galaktionov, V. A.; Posashkov, S. A.** Application of new comparison theorems to the investigation of unbounded solutions of nonlinear parabolic equations. (Russian) *Differentsial'nye Uravneniya* **22** (1986), no. 7, 1165–1173, 1285.
- [19] **Hamilton, Richard S.** Private communication.
- [20] **Herrero, M. A.; Velázquez, J. J. L.** Blow-up behaviour of one-dimensional semilinear parabolic equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **10** (1993), no. 2, 131–189.
- [21] **Herrero, Miguel A.; Velázquez, Juan J. L.** Explosion de solutions d'équations paraboliques semilinéaires supercritiques. (French) [Blowup of solutions of supercritical semilinear parabolic equations] *C. R. Acad. Sci. Paris Sér. I Math.* **319** (1994), no. 2, 141–145.
- [22] **Herrero, M. A.; Velázquez, J. J. L.** Blow-up behaviour of one-dimensional semilinear parabolic equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **10** (1993), no. 2, 131–189.
- [23] **Huisken, Gerhard; Sinestrari, Carlo.** Convexity estimates for mean curvature flow and singularities of mean convex surfaces. *Acta Math.* **183** (1999), no. 1, 45–70.
- [24] **Ilmanen, Tom; Knopf, Dan.** A lower bound for the diameter of solutions to the Ricci flow with nonzero $H^1(M^n; \mathbb{R})$. *Math. Res. Lett.* **10** (2003), no. 2, 161–168.
- [25] **King, J. R.** Self-similar behaviour for the equation of fast nonlinear diffusion. *Philos. Trans. Roy. Soc. London Ser. A* **343** (1993), 337–375.
- [26] **Mueller, Carl E.; Weissler, Fred B.** Single point blow-up for a general semilinear heat equation. *Indiana Univ. Math. J.* **34** (1985), no. 4, 881–913.
- [27] **Perelman, Grisha.** The entropy formula for the Ricci flow and its geometric applications. [arXiv:math.DG/0211159](http://arxiv.org/abs/math.DG/0211159).
- [28] **Sesum, Natasa.** Curvature tensor under the Ricci flow. [arXiv:math.DG/0311397](http://arxiv.org/abs/math.DG/0311397).
- [29] **Simon, Miles.** A class of Riemannian manifolds that pinch when evolved by Ricci flow. *Manuscripta Math.* **101** (2000), no. 1, 89–114.
- [30] **Topping, Peter.** Diameter control under Ricci flow. <http://www.maths.warwick.ac.uk/~topping/>
- [31] **van den Berg, Jan Bouwe; Hulshof, Josephus; King, John R.** Formal asymptotics of bubbling in the harmonic map heat flow. *SIAM J. Appl. Math.* **63** (2003), no. 5, 1682–1717 (electronic).

- [32] **Weissler, Fred B.** Single point blow-up for a semilinear initial value problem. *J. Differential Equations* **55** (1984), no. 2, 204–224.

(Sigurd Angenent) UNIVERSITY OF WISCONSIN - MADISON

E-mail address: `angenent@math.wisc.edu`

URL: `http://www.math.wisc.edu/~angenent/`

(Dan Knopf) UNIVERSITY OF TEXAS AT AUSTIN

E-mail address: `danknopf@math.utexas.edu`

URL: `http://www.ma.utexas.edu/~danknopf`